



Histone cluster 1 H4I (m): 293T Lysate: sc-120794

BACKGROUND

Histones are a superfamily of basic nuclear proteins that, together, are responsible for maintaining eukaryotic chromosomal structure. There are four core Histones, designated Histone H2A, Histone H2B, Histone H3 and Histone H4, all of which contribute two protein molecules that, together, form an octamer around which DNA is wrapped in repeating units known as nucleosomes. The Histone H1 subfamily of proteins interact with linker DNA between nucleosomes and are responsible for condensing chromatin into higher ordered structures. Histone cluster 1 H4I, also known as HIST1H1A, H1F1 or HIST1, is a 215 amino acid protein that localizes to the nucleus. One of several members of the Histone H1/H5 family, Histone cluster 1 H4I plays a role in the condensation of nucleosomes into higher ordered structures, thereby playing a role in transcription and, ultimately, cell cycle progression. The Histone cluster 1 H4I gene maps within a cluster of Histone genes that are found on human chromosome 6.

REFERENCES

1. Eick, S., Nicolai, M., Mumberg, D. and Doenecke, D. 1989. Human H1 histones: conserved and varied sequence elements in two H1 subtype genes. *Eur. J. Cell Biol.* 49: 110-115.
2. Albig, W., Drabent, B., Kunz, J., Kalff-Suske, M., Grzeschik, K.H. and Doenecke, D. 1993. All known human H1 histone genes except the H1⁰ gene are clustered on chromosome 6. *Genomics* 16: 649-654.
3. Lever, M.A., Th'ng, J.P., Sun, X. and Hendzel, M.J. 2000. Rapid exchange of histone H1.1 on chromatin in living human cells. *Nature* 408: 873-876.
4. Misteli, T., Gunjan, A., Hock, R., Bustin, M. and Brown, D.T. 2000. Dynamic binding of histone H1 to chromatin in living cells. *Nature* 408: 877-881.
5. Marzluff, W.F., Gongidi, P., Woods, K.R., Jin, J. and Maltais, L.J. 2002. The human and mouse replication-dependent histone genes. *Genomics* 80: 487-498.
6. Coleman, M.A., Miller, K.A., Beernink, P.T., Yoshikawa, D.M. and Albala, J.S. 2003. Identification of chromatin-related protein interactions using protein microarrays. *Proteomics* 3: 2101-2107.
7. Hendzel, M.J., Lever, M.A., Crawford, E. and Th'ng, J.P. 2004. The C-terminal domain is the primary determinant of histone H1 binding to chromatin *in vivo*. *J. Biol. Chem.* 279: 20028-20034.
8. Lin, Q., Inselman, A., Han, X., Xu, H., Zhang, W., Handel, M.A. and Skoultschi, A.I. 2004. Reductions in linker histone levels are tolerated in developing spermatocytes but cause changes in specific gene expression. *J. Biol. Chem.* 279: 23525-23535.
9. Garcia, B.A., Busby, S.A., Barber, C.M., Shabanowitz, J., Allis, C.D. and Hunt, D.F. 2004. Characterization of phosphorylation sites on histone H1 isoforms by tandem mass spectrometry. *J. Proteome Res.* 3: 1219-1227.

CHROMOSOMAL LOCATION

Genetic locus: Hist1h4i (mouse) mapping to 13 A3.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Histone cluster 1 H4I (m): 293T Lysate represents a lysate of mouse Histone cluster 1 H4I transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Histone cluster 1 H4I (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Histone cluster 1 H4I antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.