

HNF-6 (m2): 293T Lysate: sc-120850

BACKGROUND

HNF-1 (α and β), HNF-3 (α , β and γ), HNF-4 (α and γ), and HNF-6 compose, in part, a homeoprotein family designated the Hepatocyte Nuclear Factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3 α , HNF-3 β and HNF-3 γ regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 α and HNF-3 β have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 α and HNF-4 γ have been shown to be important for early embryo development. HNF-4 α is expressed in liver, kidney, pancreas, small intestine, testis and colon; and HNF-4 γ is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3 β , which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transcriptional activator for at least 22 other hepatocyte-enriched genes, including cytochrome P450 2C13 and α -1 antitrypsin.

REFERENCES

1. Bach, I., et al. 1993. More potent transcriptional activators or a trans-dominant inhibitor of the HNF1 homeoprotein family are generated by alternative RNA processing. *EMBO J.* 12: 4229-4242.
2. Kaestner, K.H., et al. 1994. The HNF-3 gene family of transcription factors in mice: gene structure, cDNA sequence, and mRNA distribution. *Genomics* 20: 377-385.
3. Drewes, T., et al. 1996. Human hepatocyte nuclear factor 4 isoforms are encoded by distinct and differentially expressed genes. *Mol. Cell. Biol.* 16: 925-931.
4. Samadani, U., et al. 1996. The transcriptional activator hepatocyte nuclear factor 6 regulates liver gene expression. *Mol. Cell. Biol.* 16: 6273-6284.
5. Pierreux, C.E., et al. 2006. The transcription factor hepatocyte nuclear factor-6 controls the development of pancreatic ducts in the mouse. *Gastroenterology* 130: 532-541.
6. Beaudry, J.B., et al. 2006. Threshold levels of hepatocyte nuclear factor 6 (HNF-6) acting in synergy with HNF-4 and PGC-1 α are required for time-specific gene expression during liver development. *Mol. Cell. Biol.* 26: 6037-6046.
7. Hatzis, P., et al. 2006. Mitogen-activated protein kinase-mediated disruption of enhancer-promoter communication inhibits hepatocyte nuclear factor 4 α expression. *Mol. Cell. Biol.* 26: 7017-7029.
8. Iyaguchi, D., et al. 2007. DNA recognition mechanism of the ONECUT homeodomain of transcription factor HNF-6. *Structure* 15: 75-83.
9. Vanhorenbeeck, V., et al. 2007. Role of the Onecut transcription factors in pancreas morphogenesis and in pancreatic and enteric endocrine differentiation. *Dev. Biol.* 305: 685-694.

CHROMOSOMAL LOCATION

Genetic locus: Onecut1 (mouse) mapping to 9 D.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

HNF-6 (m2): 293T Lysate represents a lysate of mouse HNF-6 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

HNF-6 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive HNF-6 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

HNF-6 (G-10): sc-376167 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse HNF-6 expression in HNF-6 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



HNF-6 (G-10): sc-376167. Western blot analysis of HNF-6 expression in non-transfected: sc-117752 (A) and mouse HNF-6 transfected: sc-120850 (B) 293T whole cell lysates.



HNF-6 (B-1): sc-376308. Western blot analysis of HNF-6 expression in non-transfected: sc-117752 (A) and mouse HNF-6 transfected: sc-120850 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.