

Ihh (m): 293T Lysate: sc-121020

BACKGROUND

The *Drosophila* segment polarity gene hedgehog (hh) encodes a precursor protein which undergoes autocleavage to generate amino- and carboxy-terminal peptides. Both proteins are secreted and appear to function in embryonic and imaginal disc patterning. Several vertebrate homologs of *Drosophila* hh have been identified. These include Sonic hedgehog (Shh), alternatively designated Vhh-1, Desert hedgehog (Dhh) and Indian hedgehog (Ihh). Each contain amino-terminal signal peptides and apparently function as secreted proteins involved in the mediation of various cell-cell interactions. Shh resembles *Drosophila* hh in that it is processed to generate an amino-terminal secreted peptide that is retained at or near the cell surface and a carboxy-terminal glycosylated more diffusible peptide.

REFERENCES

1. Echelard, Y., et al. 1993. Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell* 75: 1417-1430.
2. Li, W., et al. 1995. Function of protein kinase A in hedgehog signal transduction and *Drosophila* imaginal disc development. *Cell* 80: 553-562.
3. Roelink, H., et al. 1995. Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of Sonic hedgehog autoproteolysis. *Cell* 81: 445-455.
4. Fan, C.M., et al. 1995. Long-range sclerotome induction by Sonic hedgehog: direct role of the amino-terminal cleavage product and modulation by the cyclic AMP signaling pathway. *Cell* 81: 457-465.
5. Ericson, J., et al. 1995. Sonic hedgehog induces the differentiation of ventral forebrain neurons: a common signal for ventral patterning within the neural tube. *Cell* 81: 747-756.
6. Marti, E., et al. 1995. Requirement of 19K form of Sonic hedgehog for induction of distinct ventral cell types in CNS explants. *Nature* 375: 322-325.

CHROMOSOMAL LOCATION

Genetic locus: Ihh (mouse) mapping to 1 C3.

PRODUCT

Ihh (m): 293T Lysate represents a lysate of mouse Ihh transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Ihh (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Ihh antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Ihh (H-12): sc-271101 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Ihh expression in Ihh transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Ihh (H-12): sc-271101. Western blot analysis of Ihh expression in non-transfected: sc-117752 (A) and mouse Ihh transfected: sc-121020 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.