

IKK γ (m): 293T Lysate: sc-121029

BACKGROUND

The transcription factor NF κ B is retained in the cytoplasm in an inactive form by the inhibitory protein I κ B. Activation of NF κ B requires that I κ B be phospho-rylated on specific serine residues, which results in targeted degradation of I κ B. I κ B kinase α (IKK α), previously designated CHUK, interacts with I κ B α and specifically phosphorylates I κ B α on Serine 32 and 36, the sites that trigger its degradation. IKK α appears to be critical for NF κ B activation in response to proinflammatory cytokines. Phosphorylation of I κ B by IKK α is stimulated by the NF κ B inducing kinase (NIK), which itself is a central regulator for NF κ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK α , IKK β and IKK γ (also designated NEMO), and each appear to make essential contributions to I κ B phosphorylation.

REFERENCES

- Verma, I.M., Stevenson, J.K., Schwarz, E.M., Van Antwerp, D. and Miyamoto, S. 1995. Rel/NF κ B/I κ B family: intimate tales of association and dissociation. *Genes Dev.* 9: 2723-2735.
- Thanos, D. and Maniatis, T. 1995. NF κ B: a lesson in family values. *Cell* 80: 529-532.
- Connelly, M.A. and Marcu, K.B. 1995. CHUK, a new member of the helix-loop-helix and leucine zipper families of interacting proteins, contains a serine-threonine kinase catalytic domain. *Cell. Mol. Biol. Res.* 41: 537-549.
- DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. 1997. A cytokine-responsive I κ B kinase that activates the transcription factor NF κ B. *Nature* 388: 548-554.
- Régnier, C.H., Song, H.Y., Goa, X., Goeddel, D.V., Cao, Z. and Rothe, M. 1997. Identification and characterization of an I κ B kinase. *Cell* 90: 373-383.
- Malinin, N.L., Boldin, M.P., Kovalenko, A.V. and Wallach, D. 1997. MAP3K-related kinase involved in NF κ B induction by TNF, CD95 and IL-1. *Nature* 385: 540-544.
- Song, H.Y., Régnier, C.H., Kirschning, C.J., Goeddel, D.V. and Rothe, M. 1997. Tumor necrosis factor (TNF)-mediated kinase cascades: bifurcation of nuclear factor κ B and c-jun N-terminal kinase (JNK/SAPK) pathways at TNF receptor-associated factor 2. *Proc. Natl. Acad. Sci. USA* 94: 9792-9296.
- Zandi, E., Rothwarf, D.M., Delhase, M., Hayakawa, M. and Karin, M. 1997. The I κ B kinase complex (IKK) contains two kinase subunits, IKK α and IKK β , necessary for I κ B phosphorylation and NF κ B activation. *Cell* 91: 243-252.
- Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S.T., Weil, R., Agou, F., Kirk, H.E., Kay, R.J. and Israel, A. 1998. Complementation cloning of NEMO, a component of the I κ B kinase complex essential for NF κ B activation. *Cell* 93: 1231-1240.

CHROMOSOMAL LOCATION

Genetic locus: Ikbkg (mouse) mapping to X A7.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

IKK γ (m): 293T Lysate represents a lysate of mouse IKK γ transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

IKK γ (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive IKK γ antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

IKK γ (F-10): sc-166398 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse IKK γ expression in IKK γ transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



IKK γ (F-10): sc-166398. Western blot analysis of IKK γ expression in non-transfected: sc-117752 (A) and mouse IKK γ transfected: sc-121029 (B) 293T whole cell lysates.



IKK γ (F-12): sc-166397. Western blot analysis of IKK γ expression in non-transfected: sc-117752 (A) and mouse IKK γ transfected: sc-121029 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.