



MLF2 (m): 293T Lysate: sc-121684

BACKGROUND

Myeloid leukemia factor (MLF) proteins typically demonstrate highest levels of expression in testis, ovary, skeletal muscle, heart, kidney and colon tissues, and lower levels of expression in spleen, thymus and peripheral blood leukocytes. MLF proteins play a role in normal hemopoietic differentiation as well as in erythroid/myeloid lineage switching. MLF2 is a ubiquitously expressed, 248 amino acid protein which shares 40% sequence identity with myeloid leukemia factor 1 (MLF1). MLF2 maps to chromosome 12p13.31, a region that is often associated with translocations in acute leukemias of lymphoid and myeloid origin. However, no alterations in the structure of the MLF2 locus in patients shown to have 12p translocations have been discovered.

REFERENCES

1. Kuefer, M.U., Look, A.T., Williams, D.C., Valentine, V., Naeve, C.W., Behm, F.G., Mullersman, J.E., Yoneda-Kato, N., Montgomery, K., Kucherlapati, R. and Morris, S.W. 1996. cDNA cloning, tissue distribution, and chromosomal localization of myelodysplasia/myeloid leukemia factor 2 (MLF2). *Genomics* 35: 392-396.
2. Fujimura, H. 1998. Growth inhibition of *Saccharomyces cerevisiae* by the immunosuppressant leflunomide is due to the inhibition of uracil uptake via Fur4p. *Mol. Gen. Genet.* 260: 102-107.
3. Ohno, K., Takahashi, Y., Hirose, F., Inoue, Y.H., Taguchi, O., Nishida, Y., Matsukage, A. and Yamaguchi, M. 2001. Characterization of a *Drosophila* homologue of leukemia factor (MLF). *Gene* 260: 133-143.
4. Shi, Y.W., Hu, W.X. and Tang, L.J. 2003. Gene expression profile changes in human multiple myeloma. *Hunan Yi Ke Da Xue Xue Bao* 28: 201-205.
5. Noé, V., Peñuelas, S., Lamuela-Raventós, R.M., Permanyer, J., Ciudad, C.J. and Izquierdo-Pulido, M. 2004. Epicatechin and a cocoa polyphenolic extract modulate gene expression in human Caco-2 cells. *J. Nutr.* 134: 2509-2516.
6. Kim, W.Y., Fayazi, Z., Bao, X., Higgins, D. and Kazemi-Esfarjani, P. 2005. Evidence for sequestration of polyglutamine inclusions by *Drosophila* myeloid leukemia factor. *Mol. Cell. Neurosci.* 29: 536-544.

CHROMOSOMAL LOCATION

Genetic locus: Mlf2 (mouse) mapping to 6 F2.

PRODUCT

MLF2 (m): 293T Lysate represents a lysate of mouse MLF2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

MLF2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MLF2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.