

MSK2 (m): 293T Lysate: sc-121804

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including cell division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine (Ser/Thr) protein kinases. MSK2, also known as RPS6KA4 (Ribosomal Protein S6 kinase, 90 kDa, polypeptide 4) or RSKB, is a 772 amino acid protein that localizes to the nucleus and contains one AGC-kinase C-terminal domain and two protein kinase domains. Using magnesium as a cofactor, MSK2 functions as a Ser/Thr kinase that is thought to play a role in the regulation of growth-factor and stress-induced transcriptional activation, specifically by catalyzing the ATP-dependent phosphorylation of target proteins. Multiple isoforms of MSK2 exist due to alternative splicing events.

REFERENCES

1. Deak, M., et al. 1998. Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38, and may mediate activation of CREB. *EMBO J.* 17: 4426-4441.
2. Pierrat, B., et al. 1998. RSK-B, a novel ribosomal S6 kinase family member, is a CREB kinase under dominant control of p38 α mitogen-activated protein kinase (p38 α MAPK). *J. Biol. Chem.* 273: 29661-29671.
3. Online Mendelian Inheritance in Man, OMIM™. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 603606. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Gudi, T., et al. 2000. NO activation of Fos promoter elements requires nuclear translocation of G-kinase I and CREB phosphorylation but is independent of MAP kinase activation. *Oncogene* 19: 6324-6333.
5. Tomás-Zuber, M., et al. 2001. C-terminal elements control location, activation threshold, and p38 docking of ribosomal S6 kinase B (RSKB). *J. Biol. Chem.* 276: 5892-5899.
6. Prymakowska-Bosak, M., et al. 2001. Mitotic phosphorylation prevents the binding of HMGN proteins to chromatin. *Mol. Cell. Biol.* 21: 5169-5178.
7. Wiggan, G.R., et al. 2002. MSK1 and MSK2 are required for the mitogen- and stress-induced phosphorylation of CREB and ATF1 in fibroblasts. *Mol. Cell. Biol.* 22: 2871-2881.
8. Soloaga, A., et al. 2003. MSK2 and MSK1 mediate the mitogen- and stress-induced phosphorylation of histone H3 and HMG-14. *EMBO J.* 22: 2788-2797.
9. Gesser, B., et al. 2007. Dimethylfumarate specifically inhibits the mitogen and stress-activated kinases 1 and 2 (MSK1/2): possible role for its anti-psoriatic effect. *J. Invest. Dermatol.* 127: 2129-2137.

CHROMOSOMAL LOCATION

Genetic locus: Rps6ka4 (mouse) mapping to 19 A.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

MSK2 (m): 293T Lysate represents a lysate of mouse MSK2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.


APPLICATIONS

MSK2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MSK2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

MSK2 (2934C1a): sc-130653 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse MSK2 expression in MSK2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



MSK2 (2934C1a): sc-130653. Western blot analysis of MSK2 expression in non-transfected: sc-117752 (A) and mouse MSK2 transfected: sc-121804 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.