

NRK1 (m): 293T Lysate: sc-122131

BACKGROUND

Nicotinamide adenine dinucleotide (NAD⁺) is an essential cofactor involved in fundamental processes in cell metabolism. NRK1 (Nicotinamide riboside kinase 1), also known as RibosylNicotinamide kinase 1, is a 199 amino acid enzyme is involved in the synthesis of NAD⁺ through nicotinamide mononucleotide using nicotinamide riboside as the precursor. Nicotinamide riboside has been identified as a nutrient in milk, suggesting that it is a useful compound for elevating the NAD⁺ levels in humans. NRK1 also phosphorylates the anti-cancer drugs tiazofurin and 3-deazaguanosine, which converts them into toxic NAD⁺ analogs and leads to the inhibition of guanine nucleotide biosynthesis. There are two isoforms of NRK1 that are produced as a result of alternative splicing events.

REFERENCES

1. Sasiak, K. and Saunders, P.P. 1996. Purification and properties of a human nicotinamide ribonucleoside kinase. *Arch. Biochem. Biophys.* 333: 414-418.
2. Bieganski, P. and Brenner, C. 2004. Discoveries of nicotinamide riboside as a nutrient and conserved NRK genes establish a Preiss-Handler independent route to NAD⁺ in fungi and humans. *Cell* 117: 495-502.
3. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 608704. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Belenky, P., Racette, F.G., Bogan, K.L., McClure, J.M., Smith, J.S. and Brenner, C. 2007. Nicotinamide riboside promotes Sir2 silencing and extends lifespan via Nrk and Urh1/Pnp1/Meu1 pathways to NAD⁺. *Cell* 129: 473-484.
5. Ma, B., Pan, S.J., Zupancic, M.L. and Cormack, B.P. 2007. Assimilation of NAD⁺ precursors in *Candida glabrata*. *Mol. Microbiol.* 66: 14-25.
6. Tempel, W., Rabeh, W.M., Bogan, K.L., Belenky, P., Wojcik, M., Seidle, H.F., Nedyalkova, L., Yang, T., Sauve, A.A., Park, H.W. and Brenner, C. 2007. Nicotinamide riboside kinase structures reveal new pathways to NAD⁺. *PLoS Biol.* 5: e263.
7. Khan, J.A., Xiang, S. and Tong, L. 2007. Crystal structure of human nicotinamide riboside kinase. *Structure* 15: 1005-1013.
8. Belenky, P., Bogan, K.L. and Brenner, C. 2007. NAD⁺ metabolism in health and disease. *Trends Biochem. Sci.* 32: 12-19.
9. Bogan, K.L. and Brenner, C. 2008. Nicotinic acid, nicotinamide, and nicotinamide riboside: a molecular evaluation of NAD⁺ precursor vitamins in human nutrition. *Annu. Rev. Nutr.* 28: 115-130.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: BC016495 (mouse) mapping to 19 B.

PRODUCT

NRK1 (m): 293T Lysate represents a lysate of mouse NRK1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

NRK1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive NRK1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


NRK1 (F-8): sc-398852 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse NRK1 expression in NRK1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



NRK1 (F-8): sc-398852. Western blot analysis of NRK1 expression in non-transfected: sc-117752 (A) and mouse NRK1 transfected: sc-122131 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.