

Pirh2 (m): 293T Lysate: sc-122592

BACKGROUND

Pirh2, also known as androgen receptor N-terminal-interacting protein (ARNIP), ZN363 or CHIMP, has p53-induced ubiquitin-protein ligase activity, promoting p53 degradation. The protein physically interacts with p53 and the resulting degradation of p53 renders Pirh2 an oncogenic protein, as the loss of p53 function contributes to malignant tumor development. The gene encoding for the protein maps to chromosome 4q21.1; transcription of this gene is regulated by p53. Pirh2 expression decreases the level of p53, and a decrease of endogenous Pirh2 expression increases p53 levels. Pirh2 is therefore considered, together with Mdm2, to act as a negative regulator of p53 function.

REFERENCES

1. Beitel, L.K., Elhaji, Y.A., Lumbroso, R., Wing, S.S., Panet-Raymond, V., Gottlieb, B., Pinsky, L. and Trifiro, M.A. 2002. Cloning and characterization of an androgen receptor N-terminal-interacting protein with ubiquitin-protein ligase activity. *J. Mol. Endocrinol.* 29: 41-60
2. Leng, R.P., Lin, Y., Ma, W., Wu, H., Lemmers, B., Chung, S., Parant, J.M., Lozano, G., Hakem, R. and Benchimol, S. 2003. Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53 degradation. *Cell* 112: 779-791.
3. Duan, W., Gao, L., Druhan, L.J., Zhu, W.G., Morrison, C., Otterson, G.A. and Villalona-Calero, M.A. 2004. Expression of Pirh2, a newly identified ubiquitin protein ligase, in lung cancer. *J. Natl. Cancer Inst.* 96: 1718-1721.
4. Corcoran, C.A., Huang, Y. and Sheikh, M.S. 2004. The p53 paddy wagon: COP1, Pirh2 and MDM2 are found resisting apoptosis and growth arrest. *Cancer Biol. Ther.* 3: 721-725.
5. Dornan, D., Wertz, I., Shimizu, H., Arnott, D., Frantz, G.D., Dowd, P., O'Rourke, K., Koeppen, H. and Dixit, V.M. 2004. The ubiquitin ligase COP1 is a critical negative regulator of p53. *Nature* 429: 86-92.
6. Martorati, A., Doumont, G., Alcalay, M., Bellefroid, E., Pelicci, P.G. and Marine, J.C. 2005. dapk1, encoding an activator of a p19ARF-p53-mediated apoptotic checkpoint, is a transcription target of p53. *Oncogene* 24: 1461-1466.
7. Zhang, L., Li, J., Wang, C., Ma, Y. and Huo, K. 2005. A new human gene hNTKL-BP1 interacts with hPirh2. *Biochem. Biophys. Res. Commun.* 330: 293-297.
8. Dulloo, I. and Sabapathy, K. 2005. Transactivation-dependent and -independent regulation of p73 stability. *J. Biol. Chem.* 280: 28203-28214.

CHROMOSOMAL LOCATION

Genetic locus: Rchy1 (mouse) mapping to 5 E2.

PRODUCT

Pirh2 (m): 293T Lysate represents a lysate of mouse Pirh2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Pirh2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Pirh2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.