# PRX IV (m): 293T Lysate: sc-122810



The Power to Question

## **BACKGROUND**

The peroxiredoxin (PRX) family comprises six antioxidant proteins, PRX I, II, III, IV, V and VI, which protect cells from reactive oxygen species (ROS) by preventing the metal-catalyzed oxidation of enzymes. The PRX proteins primarily utilize thioredoxin as the electron donor for antioxidation, although they are fairly promiscuous with regard to the hydroperoxide substrate. In addition to protection from ROS, peroxiredoxins are also involved in cell proliferation, differentiation and gene expression. PRX I, II, IV and VI show diffuse cytoplasmic localization, while PRX III and V exhibit distinct mitochondrial localization. The human PRX I gene encodes a protein that is expressed in several tissues, including liver, kidney, testis, lung and nervous system. PRX II is expressed in testis, while PRX III shows expression in lung. PRX I, II and III are overexpressed in breast cancer and may be involved in its development or progression. Upregulated protein levels of PRX I and II in Alzheimer's disease (AD) and Down syndrome (DS) indicate the involvement of PRX I and II in their pathogenesis. The human PRX IV gene is abundantly expressed in many tissues. PRX IV exists as a precursor protein, which is only detected in testis and a processed, secreted form. PRX V also exists as two forms, designated long and short. Like PRX IV, the long form of PRX V is highly expressed in testis. The short form of PRX V is more widely expressed, with high expression in liver, kidney, heart and lung. PRX VI,  $\alpha$ 1-Cys peroxiredoxin (also known as antioxidant protein 2 or AOP2), is highly expressed in most tissues, particularly in epithelial cells. Localized to the cell cytosol, PRX VI functions independently of other peroxiredoxins and antioxidant proteins, specializing in antioxidant defense, lung phospholipid metabolism and protection of keratinocytes from cell death induced by reactive oxygen species.

# REFERENCES

- 1. Iwahara, S., et al. 1995. Purification, characterization and cloning of a hemebinding protein (23 kDa) in rat liver cytosol. Biochemistry 34: 13398-13406.
- Butterfield, L.H., et al. 1999. From cytoprotection to tumor suppression: the multifactorial role of peroxiredoxins. Antioxid. Redox Signal. 1: 385-402.
- 3. Mizusawa, H., et al. 2000. Peroxiredoxin I (macrophage 23 kDa stress protein) is highly and widely expressed in the rat nervous system. Neurosci. Lett. 283: 57-60.
- Noh, D.Y., et al. 2001. Overexpression of peroxiredoxin in human breast cancer. Anticancer Res. 21: 2085-2090.
- Kim, S.H., et al. 2001. Protein levels of human peroxiredoxin subtypes in brains of patients with Alzheimer's disease and Down syndrome. J. Neural Transm. Suppl. 223-235.
- 6. Kinnula, V.L., et al. 2002. Cell specific expression of peroxiredoxins in human lung and pulmonary sarcoidosis. Thorax 57: 157-164.
- 7. Hofmann, B., et al. 2002. Peroxiredoxins. Biol. Chem. 383: 347-364.
- 8. Fujii, J., et al. 2002. Advances in our understanding of peroxiredoxin, a multifunctional, mammalian redox protein. Redox Rep. 7: 123-130.
- 9. Wang, X., et al. 2003. Mice with targeted mutation of peroxiredoxin 6 develop normally but are susceptible to oxidative stress. J. Biol. Chem. 278: 25179-25190.

#### **CHROMOSOMAL LOCATION**

Genetic locus: Prdx4 (mouse) mapping to X F3.

#### **PRODUCT**

PRX IV (m): 293T Lysate represents a lysate of mouse PRX IV transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

#### **APPLICATIONS**

PRX IV (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PRX IV antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## **STORAGE**

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**