



STAMPB (m2): 293T Lysate: sc-123804

BACKGROUND

STMBP (STAM binding protein), also known as AMSH, is a 424 amino acid protein belonging to the peptidase M67C family. Ubiquitously expressed, STMBP functions as a zinc metalloprotease that specifically cleaves "Lys-63"-linked polyubiquitin chains. STMBP is able to oppose the ubiquitin-dependent sorting of receptors to lysosomes. STMBP may play a role in signal transduction for cell growth and Myc induction mediated by IL-2 and GM-CSF. It is suggested that STMBP potentiates BMP (bone morphogenetic protein) signaling by antagonizing the inhibitory action of Smad6 and Smad7. STMBP consists of the JAMM motif, which is essential for the protease activity, and is inhibited by N-ethylmaleimide.

REFERENCES

1. Itoh, F., et al. 2001. Promoting bone morphogenetic protein signaling through negative regulation of inhibitory Smads. *EMBO J.* 20: 4132-4142.
2. McCullough, J., et al. 2004. AMSH is an endosome-associated ubiquitin isopeptidase. *J. Cell Biol.* 166: 487-492.
3. Li, H., et al. 2004. An RNF11: Smurf2 complex mediates ubiquitination of the AMSH protein. *Oncogene* 23: 1801-1808.
4. Herrera-Vigener, F., et al. 2006. AMSH regulates calcium-sensing receptor signaling through direct interactions. *Biochem. Biophys. Res. Commun.* 347: 924-930.
5. McCullough, J., et al. 2006. Activation of the endosome-associated ubiquitin isopeptidase AMSH by STAM, a component of the multivesicular body-sorting machinery. *Curr. Biol.* 16: 160-165.
6. Nakamura, M., et al. 2006. Clathrin anchors deubiquitinating enzymes, AMSH and AMSH-like protein, on early endosomes. *Genes Cells* 11: 593-606.
7. Agromayor, M., et al. 2006. Interaction of AMSH with ESCRT-III and deubiquitination of endosomal cargo. *J. Biol. Chem.* 281: 23083-23091.
8. Ma, Y.M., et al. 2007. Targeting of AMSH to endosomes is required for epidermal growth factor receptor degradation. *J. Biol. Chem.* 282: 9805-9812.
9. Sato, Y., et al. 2008. Structural basis for specific cleavage of Lys-63-linked polyubiquitin chains. *Nature* 455: 358-362.

CHROMOSOMAL LOCATION

Genetic locus: Stambp (mouse) mapping to 6 C3.

PRODUCT

STAMPB (m2): 293T Lysate represents a lysate of mouse STAMPB transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

STAMPB (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive STAMPB antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.