



TES (m2): 293T Lysate: sc-123983

BACKGROUND

TES was originally identified as a candidate tumour suppressor gene and has been found to encode a novel focal adhesion protein called TES or Testin. TES localises to cell-cell contacts and Actin stress fibres, and interacts with a variety of cytoskeletal proteins including Zyxin, Mena, VASP, Talin and Actin. The ability of TES to associate with α -actinin, paxillin and Zyxin is dependent on the conformational state of the molecule. TES contains three LIM zinc-binding domains and may act as a tumor suppressor. Overexpression of the TES gene results in increased cell spreading and decreased cell motility.

REFERENCES

1. Garvalov, B.K., Higgins, T.E., Sutherland, J.D., Zettl, M., Scaplehorn, N., Kocher, T., Piddini, E., Griffiths, G. and Way, M. 2003. The conformational state of TES regulates its Zyxin-dependent recruitment to focal adhesions. *J. Cell Biol.* 161: 33-39.
2. Coutts, A.S., MacKenzie, E., Griffith, E. and Black, D.M. 2003. TES is a novel focal adhesion protein with a role in cell spreading. *J. Cell. Sci.* 116: 897-906.
3. Chene, L., Giroud, C., Desgrandchamps, F., Boccon-Gibod, L., Cussenot, O., Berthon, P. and Latil, A. 2004. Extensive analysis of the 7q31 region in human prostate tumors supports TES as the best candidate tumor suppressor gene. *Int. J. Cancer* 111: 798-804.
4. Griffith, E., Coutts, A.S. and Black, D.M. 2005. RNAi knockdown of the focal adhesion protein TES reveals its role in Actin stress fibre organisation. *Cell Motil. Cytoskeleton* 60: 140-152.
5. Rotter, B., Bournier, O., Nicolas, G., Dhermy, D. and Lecomte, M.C. 2005. AlphaII-spectrin interacts with TES and EVL, two Actin-binding proteins located at cell contacts. *Biochem. J.* 388: 631-638.

CHROMOSOMAL LOCATION

Genetic locus: Tes (mouse) mapping to 6 A2.

PRODUCT

TES (m2): 293T Lysate represents a lysate of mouse TES transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TES (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TES antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.