

Tim17 (m): 293T Lysate: sc-124058

BACKGROUND

Translocation of nuclear encoded preproteins into the mitochondrial matrix requires the coordinated action of the translocases Tom and Tim, which are located in the outer mitochondrial membrane and the inner membrane, respectively. The mitochondrial preprotein translocases of the outer membrane (Tom) is a multi-subunit protein that contains at least eight proteins: four import receptor subunits (Tom70, Tom37, Tom22, and Tom20), three small proteins (Tom7, Tom6, and Tom5), and a structural component of the outer membrane channel (Tom40). The Tom machinery involves the import receptors, which initiate the binding of cytosolically synthesized preproteins to the outer membrane, and a general import pore (GIP), which promotes the translocation of various pre-proteins into the mitochondria. The Tim channel imports nuclear-encoded mitochondrial preproteins, and it involves three proteins, Tim17, Tim23 and Tim44, which are represented at equimolar ratios. Tim17 is expressed as two isoforms Tim17a and Tim17b, which differ only in their C-termini sequences, and like Tim23, these proteins are ubiquitously expressed in fetal and adult tissues. Tim17 and Tim23 are integral membrane proteins that comprise the structural elements of the inner membrane channel by which the preproteins are transferred. The Tim44, on the other hand, is a largely hydrophilic protein that recruits the matrix located Hsp70 to the site where the preprotein emerges from the Tim channel.

REFERENCES

1. Neupert, W. Protein import into mitochondria. 1997. Annu. Rev. Biochem. 66: 863-917.
2. Yano, M., Kanazawa, M., Terada, K., Takeya, M., Hoogenraad, N. and Mori, M. 1998. Functional analysis of human mitochondrial receptor Tom20 for protein import into mitochondria. J. Biol. Chem. 273: 26844-26851.
3. Brix, J., Rudiger, S., Bukau, B., Schneider-Mergener, J. and Pfanner, N. 1999. Distribution of binding sequences for the mitochondrial import receptors Tom20, Tom22, and Tom70 in a presequence-carrying preprotein and a non-cleavable preprotein. J. Biol. Chem. 274: 16522-16530.
4. Bauer, M.F., Gempel, K., Reichert, A.S., Rappold, G.A., Lichtner, P., Gerbitz, K.D., Neupert, W., Brunner, M. and Hofmann, S. 1999. Genetic and structural characterization of the human mitochondrial inner membrane translocase. J. Mol. Biol. 289: 69-82.
5. Schulke, N., Sepuri, N.B., Gordon, D.M., Saxena, S., Dancis, A. and Pain, D. 1999. A multisubunit complex of outer and inner mitochondrial membrane protein translocases stabilized *in vivo* by translocation intermediates. J. Biol. Chem. 274: 22847-22854.
6. Moro, F., Sirrenberg, C., Schneider, H.C., Neupert, W. and Brunner, M. 1999. The Tim17.23 preprotein translocase of mitochondria: composition and function in protein transport into the matrix. EMBO J. 18: 3667-3675.
7. Ryan, M.T., Wagner, R. and Pfanner, N. 2000. The transport machinery for the import of preproteins across the outer mitochondrial membrane. Int. J. Biochem. Cell Biol. 32: 13-21.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Timm17a (mouse) mapping to 1 E4.

PRODUCT

Tim17 (m): 293T Lysate represents a lysate of mouse Tim17 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Tim17 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Tim17 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.