



VAP-A (m): 293T Lysate: sc-124538

BACKGROUND

SNAREs are compartmentally specific, integral membrane proteins that are involved in the fusion of membranes and the transport of intracellular proteins. SNAREs are expressed at high levels in all cell types. VAMP-associated protein A (VAP-A) is a SNARE regulator with high levels of expression in the intestine during late embryogenesis and early neonatal development. VAP-A binds to a wide range of SNAREs and fusion-related proteins, including Syntaxin 1A, rBet1, rSec22, α SNAP and NSF. This suggests that VAP-A may play a more general role in SNARE-mediated vesicle traffic between the ER and Golgi in nonpolarized cells. VAP-A also mediates traffic in cell membranes and may play an important role in modulating intestinal smooth muscle cell differentiation. VAP-A and p48 interact to form a stable complex in mammalian cells.

REFERENCES

- Butler, K.L., Sinclair, K.E., Henderson, V.J., McKinney, G., Mesidor, D.A., Katon-Benitez, I. and Weaver, W.L. 1999. The chest radiograph in critically ill surgical patients is inaccurate in predicting ventilator-associated pneumonia. *Am. Surg.* 65: 805-810.
- Nishimura, Y., Hayashi, M., Inada, H. and Tanaka, T. 1999. Molecular cloning and characterization of mammalian homologues of vesicle-associated membrane protein-associated (VAMP-associated) proteins. *Biochem. Biophys. Res. Commun.* 254: 21-26.
- Weir, M.L., Xie, H., Klip, A. and Trimble, W.S. 2001. VAP-A binds promiscuously to both v- and tSNAREs. *Biochem. Biophys. Res. Commun.* 286: 616-621.
- Wyles, J.P., McMaster, C.R. and Ridgway, N.D. 2002. Vesicle-associated membrane protein-associated protein A (VAP-A) interacts with the oxysterol-binding protein to modify export from the endoplasmic reticulum. *J. Biol. Chem.* 277: 29908-29918.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605703. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Gabetta, V., Trzyna, W., Phiel, C. and McHugh, K.M. 2003. Vesicle-associated protein A is differentially expressed during intestinal smooth muscle cell differentiation. *Dev. Dyn.* 228: 11-20.
- Ettayebi, K. and Hardy, M.E. 2003. Norwalk virus nonstructural protein p48 forms a complex with the SNARE regulator VAP-A and prevents cell surface expression of vesicular stomatitis virus G protein. *J. Virol.* 77: 11790-11797.
- Jordan, M.C., Harrington, J.R., Cohen, N.D., Tsois, R.M., Dangott, L.J., Weinberg, E.D. and Martens, R.J. 2003. Effects of iron modulation on growth and viability of *Rhodococcus equi* and expression of virulence-associated protein A. *Am. J. Vet. Res.* 64: 1337-1346.
- Wyles, J.P. and Ridgway, N.D. 2004. VAMP-associated protein A regulates partitioning of oxysterol-binding protein-related protein-9 between the endoplasmic reticulum and Golgi apparatus. *Exp. Cell Res.* 297: 533-547.

CHROMOSOMAL LOCATION

Genetic locus: Vapa (mouse) mapping to 17 E1.2.

PRODUCT

VAP-A (m): 293T Lysate represents a lysate of mouse VAP-A transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

VAP-A (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive VAP-A antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.