



ZNF277 (m): 293T Lysate: sc-124784

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF277 (zinc finger protein 277), also known as NRIF4 or ZNF277P, is a 438 amino acid protein that contains two C₂H₂-type zinc fingers and is thought to localize to the nucleus. Expressed in fetal liver, lymph node, spleen and peripheral blood leukocytes, ZNF277 may function as a transcriptional repressor that plays a role in cell growth and differentiation.

REFERENCES

1. Liang, H., Guo, W. and Nagarajan, L. 2000. Chromosomal mapping and genomic organization of an evolutionarily conserved zinc-finger gene ZNF277. *Genomics* 66: 226-228.
2. Rousseau-Merck, M.F., Koczan, D., Legrand, I., Möller, S., Autran, S. and Thiesen, H.J. 2002. The KIX zinc-finger genes: genome wide mapping of 368 ZNF PAC clones with zinc-finger gene clusters predominantly in 23 chromosomal loci are confirmed by human sequences annotated in Ensembl. *Cytogenet. Genome Res.* 98: 147-153.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605465. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Nakamura, M., Runko, A.P. and Sagerström, C.G. 2004. A novel subfamily of zinc-finger genes involved in embryonic development. *J. Cell. Biochem.* 93: 887-895.
5. Englbrecht, C.C., Schoof, H. and Böhm, S. 2004. Conservation, diversification and expansion of C₂H₂ zinc-finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39.
6. O'Geen, H., Squazzo, S.L., Iyengar, S., Blahnik, K., Rinn, J.L., Chang, H.Y., Green, R. and Farnham, P.J. 2007. Genome-wide analysis of KAP1 binding suggests autoregulation of KRAB-ZNFs. *PLoS Genet.* 3: e89.

CHROMOSOMAL LOCATION

Genetic locus: Zfp277 (mouse) mapping to 12 B1.

PRODUCT

ZNF277 (m): 293T Lysate represents a lysate of mouse ZNF277 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

ZNF277 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ZNF277 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.