

MEK kinase-3 (m): 293T Lysate: sc-125599

BACKGROUND

Mitogen-activated protein (MAP) kinase cascades are activated by various extracellular stimuli including growth factors. The MEK kinases (also designated MAP kinase kinase kinases, MKKKs, MAP3Ks or MEKKs) phosphorylate and thereby activate the MEKs (also called MAP kinase kinases or MKKs), including ERK, JNK and p38. These activated MEKs in turn phosphorylate and activate the MAP kinases. The MEK kinases include Raf-1, Raf-B, Mos, MEK kinase-1, MEK kinase-2, MEK kinase-3, MEK kinase-4, ASK 1 (MEK kinase-5) and MAP3K6 (MEK kinase-6). MEK kinase-1 has been shown to phosphorylate MEK-1 via a Raf-independent pathway. Evidence suggests that MEK-3 is preferentially activated by MEK kinase-3 and that MEK-4 is activated by both MEK kinase-2 and MEK kinase-3. MEK kinase-4 has been shown to specifically activate the JNK pathway. ASK 1 activates both MEK-4 and MEK-3/MEK-6 pathways.

REFERENCES

1. Lange-Carter, C.A., et al. 1993. A divergence in the MAP kinase regulatory network defined by MEK kinase and Raf. *Science* 260: 315-319.
2. Guan, K.L. 1994. The mitogen activated protein kinase signal transduction pathway: from the cell surface to the nucleus. *Cell. Signal.* 6: 581-589.
3. Wang, X.S., et al. 1996. Molecular cloning and characterization of a novel protein kinase with a catalytic domain homologous to mitogen-activated protein kinase kinase kinase. *J. Biol. Chem.* 271: 31607-31611.
4. Fanger, G.R., et al. 1997. MEK kinases are regulated by EGF and selectively interact with Rac/Cdc42. *EMBO J.* 16: 4961-4972.
5. Gerwins, P., et al. 1997. Cloning of a novel mitogen-activated protein kinase kinase kinase, MEKK-4, that selectively regulates the c-Jun amino terminal kinase pathway. *J. Biol. Chem.* 272: 8288-8295.
6. Deacon, K., et al. 1997. Characterization of the mitogen-activated protein kinase kinase-4 (MKK4)/c-Jun NH₂-terminal kinase-1 and MKK3/p38 pathways regulated by MEK kinases-2 and -3. MEK kinase-3 activates MKK3 but does not cause activation of p38 kinase *in vivo*. *J. Biol. Chem.* 272: 14489-14496.

CHROMOSOMAL LOCATION

Genetic locus: Map3k3 (mouse) mapping to 11 E1.

PRODUCT

MEK kinase-3 (m): 293T Lysate represents a lysate of mouse MEK kinase-3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

MEK kinase-3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MEK kinase-3 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.