

PPIH (m): 293T Lysate: sc-125851

BACKGROUND

PPIH (peptidylprolyl isomerase H (Cyclophilin H)), also known as PPlase H, Rotamase H, CypH, CYP20 or USA-CYP (U-snRNP-associated cyclophilin SnuCyp-20), is a 177 amino acid protein that belongs to the cyclophilin-type PPlase family. PPIH may accelerate the folding of proteins and catalyzes the *cis-trans* isomerization of proline imidic peptide bonds in oligopeptides. PPIH is thought to participate in pre-mRNA splicing with processing factors PRPF3, PRPF4 and PRPF18 and may be involved in the assembly of the U4/U5/U6 tri-snRNP complex. Considered a protein chaperone, PPIH possesses PPlase activity and mediates the interactions between different proteins inside the spliceosome. PPIH contains one PPlase cyclophilin-type domain and is inhibited by Cyclosporin A.

REFERENCES

1. Chambraud, B., Rouvière-Fourmy, N., Radanyi, C., Hsiao, K., Peattie, D.A., Livingston, D.J. and Baulieu, E.E. 1993. Overexpression of p59-HBI (FKBP59), full length and domains, and characterization of PPlase activity. *Biochem. Biophys. Res. Commun.* 196: 160-166.
2. Schmidt, B., Tradler, T., Rahfeld, J.U., Ludwig, B., Jain, B., Mann, K., Rücknagel, K.P., Janowski, B., Schierhorn, A., Küllertz, G., Hacker, J. and Fischer, G. 1996. A cyclophilin-like peptidyl-prolyl *cis/trans* isomerase from *Legionella pneumophila*—characterization, molecular cloning and overexpression. *Mol. Microbiol.* 21: 1147-1160.
3. Horowitz, D.S., Kobayashi, R. and Krainer, A.R. 1997. A new cyclophilin and the human homologues of yeast Prp3 and Prp4 form a complex associated with U4/U6 snRNPs. *RNA* 3: 1374-1387.
4. Teigelkamp, S., Achsel, T., Mundt, C., Göthel, S.F., Cronshagen, U., Lane, W.S., Marahiel, M. and Lührmann, R. 1998. The 20 kD protein of human [U4/U6.U5] tri-snRNPs is a novel cyclophilin that forms a complex with the U4/U6-specific 60 kD and 90 kD proteins. *RNA* 4: 127-141.
5. Reidt, U., Wahl, M.C., Fasshauer, D., Horowitz, D.S., Lührmann, R. and Ficner, R. 2003. Crystal structure of a complex between human spliceosomal Cyclophilin H and a U4/U6 snRNP-60 K peptide. *J. Mol. Biol.* 331: 45-56.
6. Ingelfinger, D., Göthel, S.F., Marahiel, M.A., Reidt, U., Ficner, R., Lührmann, R. and Achsel, T. 2003. Two protein-protein interaction sites on the spliceosome-associated human Cyclophilin CypH. *Nucleic Acids Res.* 31: 4791-4796.
7. Galat, A. 2004. Function-dependent clustering of orthologues and paralogues of cyclophilins. *Proteins* 56: 808-820.
8. Gonzalez-Santos, J.M., Cao, H., Duan, R.C. and Hu, J. 2008. Mutation in the splicing factor Hprp3p linked to retinitis pigmentosa impairs interactions within the U4/U6 snRNP complex. *Hum. Mol. Genet.* 17: 225-239.
9. Patel, S.B. and Bellini, M. 2008. The assembly of a spliceosomal small nuclear ribonucleoprotein particle. *Nucleic Acids Res.* 36: 6482-6493.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Ppih (mouse) mapping to 4 D2.1.

PRODUCT

PPIH (m): 293T Lysate represents a lysate of mouse PPIH transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

PPIH (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PPIH antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


PPIH (C-1): sc-377217 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PPIH expression in PPIH transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PPIH (C-1): sc-377217. Western blot analysis of PPIH expression in non-transfected: sc-117752 (A) and mouse PPIH transfected: sc-125851 (B) 293T whole cell lysates.



PPIH (G-6): sc-377420. Western blot analysis of PPIH expression in non-transfected: sc-117752 (A) and mouse PPIH transfected: sc-125851 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.