



PSAP (m): 293T Lysate: sc-125862

BACKGROUND

Puromycin-sensitive aminopeptidase (PSA or PSAP) is a 100 kDa zinc metalloproteinase which degrades neuropeptides by removing amino acid residues from the amino-terminus. The protein is the most abundant aminopeptidase in the brain, however it is not exclusive to that organ. It is localized primarily in the cytoplasm, and plays a role in the metabolism of neuropeptides in nerve terminals and synaptic clefts. The human PSA gene maps to chromosome 17q21.32.

REFERENCES

1. Hui, M., et al. 1995. Changes in puromycin-sensitive aminopeptidases in postmortem schizophrenic brain regions. *Neurochem. Int.* 27: 433-441.
2. Constam, D.B., et al. 1995. Puromycin-sensitive aminopeptidase. Sequence analysis, expression, and functional characterization. *J. Biol. Chem.* 270: 26931-26939.
3. Tobler, A.R., et al. 1997. Cloning of the human puromycin-sensitive aminopeptidase and evidence for expression in neurons. *J. Neurochem.* 68: 889-897.
4. Bauer, W.O., et al. 2001. Human puromycin-sensitive aminopeptidase: cloning of 3' UTR, evidence for a polymorphism at a.a. 140 and refined chromosomal localization to 17q21. *Cytogenet. Cell Genet.* 92: 221-224.
5. Yamamoto, M., et al. 2002. Axonal transport of puromycin-sensitive aminopeptidase in rat sciatic nerves. *Neurosci. Res.* 42: 133-140.
6. Kakuta, H., et al. 2003. Fluorescent bioprobes for visualization of puromycin-sensitive aminopeptidase in living cells. *Bioorg. Med. Chem. Lett.* 13: 83-86.
7. Ma, Z., et al. 2003. Proteolytic cleavage of the puromycin-sensitive aminopeptidase generates a substrate binding domain. *Arch. Biochem. Biophys.* 415: 80-86.
8. Thompson, M.W., et al. 2003. Analysis of conserved residues of the human puromycin-sensitive aminopeptidase. *Peptides* 24: 1359-1365.
9. SWISS-PROT/TrEMBL (P55786). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: Npepps (mouse) mapping to 11 D.

PRODUCT

PSAP (m): 293T Lysate represents a lysate of mouse PSAP transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

PSAP (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PSAP antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.