

TMC6 (m2): 293T Lysate: sc-126110

BACKGROUND

TMC6 (transmembrane channel-like 6), also known as EVER1, EVIN1, EV1 or LAK-4P, is an 805 amino acid multi-pass membrane protein that localizes to the endoplasmic reticulum and belongs to the transmembrane channel family. Expressed in testis, placenta and prostate, TMC6 exists as multiple alternatively spliced isoforms and, when defective, is associated with the pathogenesis of epidermodysplasia verruciformis (EV). EV is an autosomal recessive dermatosis that is characterized by an increased susceptibility to human papillomaviruses (HPVs) and an increased rate of squamous cell carcinoma in UV-exposed skin. The gene encoding TMC6 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

1. Ramoz, N., Rueda, L.A., Bouadjar, B., Favre, M. and Orth, G. 1999. A susceptibility locus for epidermodysplasia verruciformis, an abnormal predisposition to infection with the oncogenic human papillomavirus type 5, maps to chromosome 17qter in a region containing a psoriasis locus. *J. Invest. Dermatol.* 112: 259-263.
2. Ramoz, N., Taïeb, A., Rueda, L.A., Montoya, L.S., Bouadjar, B., Favre, M. and Orth, G. 2000. Evidence for a nonallelic heterogeneity of epidermodysplasia verruciformis with two susceptibility loci mapped to chromosome regions 2p21-p24 and 17q25. *J. Invest. Dermatol.* 114: 1148-1153.
3. Keresztes, G., Mutai, H. and Heller, S. 2003. TMC and EVER genes belong to a larger novel family, the TMC gene family encoding transmembrane proteins. *BMC Genomics* 4: 24.
4. Kurima, K., Yang, Y., Sorber, K. and Griffith, A.J. 2003. Characterization of the transmembrane channel-like (TMC) gene family: functional clues from hearing loss and epidermodysplasia verruciformis. *Genomics* 82: 300-308.
5. Tate, G., Suzuki, T., Kishimoto, K. and Mitsuya, T. 2004. Novel mutations of EVER1/TMC6 gene in a Japanese patient with epidermodysplasia verruciformis. *J. Hum. Genet.* 49: 223-225.
6. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 605828. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Donfack, J., Buchinsky, F.J., Derkay, C.S., Steinberg, B.M., Choi, S.S., Conley, S.F., Meyer, C.M., McClay, J.E., Campisi, P., Hu, F.Z., Preston, R.A., Abramson, A.L., Ehrlich, G.D. and Post, J.C. 2006. Four mutations in Epidermodysplasia verruciformis 1 (EVER1) gene are not contributors to susceptibility in RRP. *Int. J. Pediatr. Otorhinolaryngol.* 70: 1235-1240.
8. Lazarczyk, M., Pons, C., Mendoza, J.A., Cassonnet, P., Jacob, Y. and Favre, M. 2008. Regulation of cellular zinc balance as a potential mechanism of EVER-mediated protection against pathogenesis by cutaneous oncogenic human papillomaviruses. *J. Exp. Med.* 205: 35-42.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Tmc6 (mouse) mapping to 11 E2.

PRODUCT

TMC6 (m2): 293T Lysate represents a lysate of mouse TMC6 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TMC6 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TMC6 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.