

VAT1 (m2): 293T Lysate: sc-126211

BACKGROUND

The storage and release of neurotransmitters in the nerve terminal is regulated by synaptic vesicles. In response to an intracellular increase in calcium levels, synaptic vesicles open and release neurotransmitters across the neuronal synapse, thereby propagating nerve impulses between neurons. VAT1 (vesicle amine transport protein 1) is a 393 amino acid integral membrane protein that is located within cholinergic synaptic vesicles. Expressed in tissues throughout the body, VAT1 belongs to the quinone oxidoreductase subfamily of zinc-containing alcohol dehydrogenase proteins and is thought to play a role in vesicular transport. Defects in the gene encoding VAT1 may be associated with endocrine disorders and tumorigenesis.

REFERENCES

1. Linial, M., Miller, K. and Scheller, R.H. 1989. VAT-1: an abundant membrane protein from torpedo cholinergic synaptic vesicles. *Neuron* 2: 1265-1273.
2. Peter, D., Finn, J.P., Klisak, I., Liu, Y., Kojis, T., Heinzmann, C., Roghani, A., Sparkes, R.S. and Edwards, R.H. 1993. Chromosomal localization of the human vesicular amine transporter genes. *Genomics* 18: 720-723.
3. Friedman, L.S., Ostermeyer, E.A., Lynch, E.D., Szabo, C.I., Anderson, L.A., Dowd, P., Lee, M.K., Rowell, S.E., Boyd, J. and King, M.C. 1994. The search for BRCA1. *Cancer Res.* 54: 6374-6382.
4. Miki, Y., Swensen, J., Shattuck-Eidens, D., Futreal, P.A., Harshman, K., Tavtigian, S., Liu, Q., Cochran, C., Bennett, L.M. and Ding, W. 1994. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 266: 66-71.
5. Smith, T.M., Lee, M.K., Szabo, C.I., Jerome, N., McEuen, M., Taylor, M., Hood, L. and King, M.C. 1996. Complete genomic sequence and analysis of 117 kb of human DNA containing the gene BRCA1. *Genome Res.* 6: 1029-1049.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604631. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Holt, M., Riedel, D., Stein, A., Schuette, C. and Jahn, R. 2008. Synaptic vesicles are constitutively active fusion machines that function independently of Ca^{2+} . *Curr. Biol.* 18: 715-722.

CHROMOSOMAL LOCATION

Genetic locus: Vat1 (mouse) mapping to 11 D.

PRODUCT

VAT1 (m2): 293T Lysate represents a lysate of mouse VAT1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

VAT1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive VAT1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.