

VHR (m): 293T Lysate: sc-126215

BACKGROUND

Mitogen-activated protein (MAP) kinases are a large class of proteins involved in signal transduction pathways that are activated by a range of stimuli and mediate a number of physiological and pathological changes in the cell. Dual specificity phosphatases (DSPs) are a subclass of the protein tyrosine phosphatase (PTP) gene superfamily, which are selective for dephosphorylating critical phosphothreonine and phosphotyrosine residues within MAP kinases. DSP gene expression is induced by a host of growth factors and/or cellular stresses, thereby negatively regulating MAP kinase superfamily members including MAPK/ERK, SAPK/JNK and p38. The members of the dual-specificity phosphatase protein family include MKP-1/CL100 (3CH134), VHR, PAC1, MKP-2, hVH-3 (B23), hVH-5, MKP-3, MKP-X, and MKP-4. Human VHR maps to chromosome 17q21.31 and encodes a 185 amino acid protein that elicits protein-serine and tyrosine phosphatase activity and is expressed in breast and ovarian tissues.

REFERENCES

1. Keyse, S.M. 1995. An emerging family of dual specificity MAP kinase phosphatases. *Biochim. Biophys. Acta* 1265: 152-160.
2. Muda, M., et al. 1997. Molecular cloning and functional characterization of a novel mitogen-activated protein kinase phosphatase, MKP-4. *J. Biol. Chem.* 272: 5141-5151.
3. Sun, H. 1998. Functional studies of dual-specificity phosphatases. *Methods Mol. Biol.* 84: 307-318.
4. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 603068. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Camps, M., et al. 2000. Dual specificity phosphatases: a gene family for control of MAP kinase function. *FASEB J.* 14: 6-16.

CHROMOSOMAL LOCATION

Genetic locus: Dusp3 (mouse) mapping to 11 D.

PRODUCT

VHR (m): 293T Lysate represents a lysate of mouse VHR transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

VHR (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive VHR antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

VHR (24): sc-136118 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse VHR expression in VHR transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



VHR (24): sc-136118. Western blot analysis of VHR expression in non-transfected: sc-117752 (A) and mouse VHR transfected: sc-126215 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.