

CENP-B (m2): 293T Lysate: sc-126624

BACKGROUND

A replicated chromosome includes two kinetochores that control chromosome segregation during mitosis. Both centromere proteins CENP-B and CENP-H are contained in the centromeric heterochromatin between kinetochores, and are involved in maintaining sister chromatid cohesion. The highly dispersed CENP-B promotes and maintains the joining of DNA satellites in the centromere. CENP-B targets centromeric α -DNA and protects it from digestion by nucleases as well as preventing DNase or restriction enzyme digestion from affecting the morphology of centromeres. CENP-H contains a coiled-coil structure and a nuclear localization signal. CENP-H is specifically and constitutively localized to kinetochores and plays a role in the organization and function of kinetochores throughout the cell cycle.

REFERENCES

1. Rieder, C.L., et al. 1998. The vertebrate cell kinetochore and its roles during mitosis. *Trends Cell Biol.* 8: 310-318.
2. Barbosa-Cisneros, O., et al. 1998. Localization of the centromere protein CENP-B using scleroderma sera and evidence for a role in centromere survival. *Rev. Rhum. Engl. Ed* 65: 15-20.
3. Sugata, N., et al. 1999. Characterization of a novel kinetochore protein, CENP-H. *J. Biol. Chem.* 274: 27343-27346.
4. Choo, K.H. 2000. Centromerization. *Trends Cell Biol.* 10: 182-188.
5. Cooke, C.A., et al. 1990. CENP-B: a major human centromere protein located beneath the kinetochore. *J. Cell Biol.* 110: 1475-1488.
6. Ohzeki, J., et al. 2002. CENP-B box is required for *de novo* centromere chromatin assembly on human alphoid DNA. *J. Cell Biol.* 159: 765-775.
7. Suzuki, N., et al. 2004. CENP-B interacts with CENP-C domains containing MIF2 regions responsible for centromere localization. *J. Biol. Chem.* 279: 5934-5946.
8. Lomonte, P., et al. 2007. Centromeric protein CENP-B proteasomal degradation induced by the viral protein ICPO. *FEBS Lett.* 581: 658-662.
9. Dolnik, A.V., et al. 2007. Dynamics of satellite binding protein CENP-B and telomere binding protein TRF2/MTBP in the nuclei of mouse spermatogenic line. *Cell Biol. Int.* 31: 316-329.

CHROMOSOMAL LOCATION

Genetic locus: Cenpb (mouse) mapping to 2 F1.

PRODUCT

CENP-B (m2): 293T Lysate represents a lysate of mouse CENP-B transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CENP-B (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CENP-B antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CENP-B (2D-7): sc-32285 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse CENP-B expression in CENP-B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™]
Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CENP-B (2D-7): sc-32285. Western blot analysis of CENP-B expression in non-transfected: sc-117752 (A) and mouse CENP-B transfected: sc-126624 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.