



FAM110A (m4): 293T Lysate: sc-126828

BACKGROUND

FAM110A (family with sequence similarity 110, member A) is a 295 amino acid protein that localizes to the cytoplasm, the cytoskeleton and the centrosome. Specifically, it co-localizes with microtubules during interphase and at spindle poles in mitosis. Expressed in thyroid, lymph node, trachea, adrenal gland, bone marrow, spleen, thymus, prostate and peripheral blood leukocyte, FAM110A is thought to interact with CSPP and play a role in proper cell cycle progression. FAM110A expression is increased during S phase and G₂/M phase. The gene encoding FAM110A maps to human chromosome 20, which houses over 60 genes and comprises approximately 2% of the human genome. Chromosome 20 contains a region with numerous genes expressed in the epididymis which are thought important for seminal production and some viewed as potential targets for male contraception.

REFERENCES

1. Prusiner, S.B. 1998. The prion diseases. *Brain Pathol.* 8: 499-513.
2. Collins, S., et al. 2001. Gerstmann-Sträussler-Scheinker syndrome, fatal familial insomnia and kuru: a review of these less common human transmissible spongiform encephalopathies. *J. Clin. Neurosci.* 8: 387-397.
3. Masullo, C. and Macchi, G. 2001. Does PRNP gene control the clinical and pathological phenotype of human spongiform transmissible encephalopathies? *Clin. Neuropathol.* 20: 19-25.
4. Joó, J.G., et al. 2006. Trisomy 20 mosaicism and nonmosaic trisomy 20: a report of 2 cases. *J. Reprod. Med.* 51: 209-212.
5. Ville, D., et al. 2006. Early pattern of epilepsy in the ring chromosome 20 syndrome. *Epilepsia* 47: 543-549.
6. Elghezal, H., et al. 2007. Ring chromosome 20 syndrome without deletions of the subtelomeric and CHRNA4-KCNQ2 genes loci. *Eur. J. Med. Genet.* 50: 441-445.
7. Kazantsev, A.G. 2007. Cellular pathways leading to neuronal dysfunction and degeneration. *Drug News Perspect.* 20: 501-509.
8. Lundwall, A. 2007. A locus on chromosome 20 encompassing genes that are highly expressed in the epididymis. *Asian J. Androl.* 9: 540-544.
9. O'Rand, M.G., et al. 2007. Eppin: an epididymal protease inhibitor and a target for male contraception. *Soc. Reprod. Fert. Suppl.* 63: 445-453.

CHROMOSOMAL LOCATION

Genetic locus: Fam110a (mouse) mapping to 2 G3.

PRODUCT

FAM110A (m4): 293T Lysate represents a lysate of mouse FAM110A transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

FAM110A (m4): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive FAM110A antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

FAM110A (B-8): sc-374558 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse FAM110A expression in FAM110A transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



FAM110A (B-8): sc-374558. Western blot analysis of FAM110A expression in non-transfected: sc-117752 (A) and mouse FAM110A transfected: sc-126828 (B) 293T whole cell lysates.



FAM110A (F-4): sc-390894. Western blot analysis of FAM110A expression in non-transfected: sc-117752 (A) and mouse FAM110A transfected: sc-126828 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.