

NMT2 (m): 293T Lysate: sc-127232

BACKGROUND

N-terminal myristoylation is a cotranslational lipid modification, which is crucial for the targeting and function of many signaling proteins. The N-myristoyltransferases (NMT 1 and NMT 2), also known as glycylpeptide N-tetradecan-oyltransferases, are cytoplasmic proteins that belong to the NMT family of proteins. The proteins in this family catalyze the addition of a myristoyl group to the N-terminal glycine residue of eukaryotic, fungal and viral proteins. They are primarily detected in heart, gut, kidney, liver and placenta.

REFERENCES

1. McIlhinney, R.A., Patel, P.B. and McGlone, K. 1994. Characterization of a polyhistidine-tagged form of human myristoyl-CoA: protein N-myristoyltransferase produced in *Escherichia coli*. *Eur. J. Biochem.* 222: 137-146.
2. Weston, S.A., Camble, R., Colls, J., Rosenbrock, G., Taylor, I., Egerton, M., Tucker, A.D., Tunncliffe, A., Mistry, A., Mancina, F., de la Fortelle, E., Irwin, J., Bricogne, G. and Paupit, R.A. 1998. Crystal structure of the anti-fungal target N-myristoyl transferase. *Nat. Struct. Biol.* 5: 213-221.
3. Rajala, R.V., Kakkar, R., Kanthan, R., Radhi, J.M., Wang, X., Wang, R., Datla, R.S. and Sharma, R.K. 2002. Altered expression and localization of N-myristoyltransferase in experimentally induced rat model of ischemia-reperfusion. *J. Cell. Biochem.* 86: 509-519.
4. Selvakumar, P., Lakshmikuttyamma, A., Lawman, Z., Bonham, K., Dimmock, J.R. and Sharma, R.K. 2004. Expression of methionine aminopeptidase 2, N-myristoyltransferase, and N-myristoyltransferase inhibitor protein 71 in HT29. *Biochem. Biophys. Res. Commun.* 322: 1012-1017.
5. Sharma, R.K. 2004. Potential role of N-myristoyltransferase in pathogenic conditions. *Can. J. Physiol. Pharmacol.* 82: 849-859.
6. Lu, Y., Selvakumar, P., Ali, K., Shrivastav, A., Bajaj, G., Resch, L., Griebel, R., Fournay, D., Meguro, K. and Sharma, R.K. 2005. Expression of N-myristoyltransferase in human brain tumors. *Neurochem. Res.* 30: 9-13.
7. Yang, S.H., Shrivastav, A., Kosinski, C., Sharma, R.K., Chen, M.H., Berthiaume, L.G., Peters, L.L., Chuang, P.T., Young, S.G. and Bergo, M.O. 2005. N-myristoyltransferase 1 is essential in early mouse development. *J. Biol. Chem.* 280: 18990-18995.
8. Price, H.P., Panethymitaki, C., Goulding, D. and Smith, D.F. 2005. Functional analysis of TbARL1, an N-myristoylated Golgi protein essential for viability in bloodstream trypanosomes. *J. Cell Sci.* 118: 831-841.
9. Pasha, M.K., Sharma, R.K. and Rajput, A.H. 2005. Increased myocardial N-myristoyltransferase activity in rotenone model of Parkinsonism. *Int. J. Mol. Med.* 15: 987-991.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Nmt2 (mouse) mapping to 2 A1.

PRODUCT

NMT2 (m): 293T Lysate represents a lysate of mouse NMT2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

NMT2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive NMT2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.