

CD14 (h3): 293 Lysate: sc-128274

BACKGROUND

Lipopolysaccharide (LPS) elicits the secretion of mediators and cytokines produced by activated macrophages and monocytes. CD14 is a glycosylphosphatidylinositol (GPI)-anchored protein found on the surfaces of monocytes and polymorphonuclear leukocytes. CD14 functions as a receptor for LPS, resulting in the secretion of various proteins. An important component in the LPS activation of monocytes through the CD14 receptor is the "adapter molecule", lipopolysaccharide binding protein (LBP). There are two forms of CD14, a membrane-associated form (mCD14), and a soluble form (sCD14). mCD14 responds to LPS alone and facilitates the secretion of proteins, while cells not expressing mCD14 fail to respond to LPS. The cells that lack mCD14 respond to LPS/LBP in the presence of sCD14.

REFERENCES

1. Simmons, D.L., Tan, S., Tenen, D.G., Nicholson-Weller, A. and Seed, B. 1989. Monocyte antigen CD14 is a phospholipid anchored membrane protein. *Blood* 73: 284-289.
2. Schumann, R.R. 1992. Function of lipopolysaccharide (LPS)-binding protein (LBP) and CD14, the receptor for LPS/LBP complexes: a short review. *Res. Immunol.* 143: 11-15.
3. Kielan, T.L. and Blecha, F. 1995. CD14 and other recognition molecules for lipopolysaccharide: a review. *Immunopharmacology* 29: 187-205.
4. Camussi, G., Mariano, F., Biancone, L., De Martino, A., Bussolati, B., Montrucchio, G. and Tobias, P.S. 1995. Lipopolysaccharide binding protein and CD14 modulate the synthesis of platelet-activating factor by human monocytes and mesangial and endothelial cells stimulated with lipopolysaccharide. *J. Immunol.* 155: 316-324.
5. Detmers, P.A., Zhou, D., Powell, D., Lichenstein, H., Kelley, M. and Pironkova, R. 1995. Endotoxin receptors (CD14) are found with CD16 (Fc γ RIII) in an intracellular compartment of neutrophils that contains alkaline phosphatase. *J. Immunol.* 155: 2085-2095.
6. Parsons, P.E., Gillespie, M.M., Moore, E.E., Moore, F.A. and Worthen, G.S. 1995. Neutrophil response to endotoxin in the adult respiratory distress syndrome: role of CD14. *Am. J. Respir. Cell Mol. Biol.* 13: 152-160.
7. Bufler, P., Stiegler, G., Schuchmann, M., Hess, S., Kruger, C., Stelter, F., Eckerskorn, C., Schutt, C. and Engelmann, H. 1995. Soluble lipopolysaccharide receptor (CD14) is released via two different mechanisms from human monocytes and CD14 transfectants. *Eur. J. Immunol.* 25: 604-610.

CHROMOSOMAL LOCATION

Genetic locus: CD14 (human) mapping to 5q31.3.

PRODUCT

CD14 (h3): 293 Lysate represents a lysate of human CD14 transfected 293 cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

CD14 (h3): 293 Lysate is suitable as a Western Blotting positive control for human reactive CD14 antibodies. Recommended use: 10-20 μ l per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

CD14 (5A3B11B5): sc-58951 is recommended as a positive control antibody for Western Blot analysis of enhanced human CD14 expression in CD14 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™
Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CD14 (5A3B11B5): sc-58951. Western blot analysis of CD14 expression in non-transfected: sc-110760 (A) and human CD14 transfected: sc-128274 (B) 293 whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.