



Factor V (h3): 293T Lysate: sc-128560

BACKGROUND

Hemostasis following tissue injury involves the deployment of essential plasma procoagulants (Prothrombin and Factors X, IX, V and VIII), which are involved in a blood coagulation cascade that leads to the formation of insoluble Fibrin clots and the promotion of platelet aggregation. Coagulation Factor V (Factor V, FV, proaccelerin, labile factor) is a 2,196 amino acid, single chain glycoprotein that is cleaved by Thrombin to yield an active, Ca^{2+} -dependent dimer. This heterodimer is essential to the blood coagulation cascade. Together with catalytic Factor Xa and Ca^{2+} on the surface of platelets or endothelial cells, Factor Va coordinates in a prothrombinase complex, which mediates proteolysis of Prothrombin into active Thrombin. Due to both the procoagulant properties of Factor V in coordinating proteolytic activation of Thrombin, and anticoagulant properties as a cofactor to activated protein C (APC), which selectively destroys FVa and FXa, alterations at the Factor V locus can contribute to hemorrhagic diathesis or thrombosis, respectively.

REFERENCES

1. Davie, E.W. and Fujikawa, K. 1975. Basic mechanisms in blood coagulation. *Annu. Rev. Biochem.* 44: 799-829.
2. Kane, W.H. and Davie, E.W. 1986. Cloning of a cDNA coding for human Factor V, a blood coagulation factor homologous to Factor VIII and ceruloplasmin. *Proc. Natl. Acad. Sci. USA* 83: 6800-6804.
3. Jenny, R.J., Pittman, D.D., Toole, J.J., Kriz, R.W., Aldape, R.A., Hewick, R.M., Kaufman, R.J. and Mann, K.G. 1987. Complete cDNA and derived amino acid sequence of human Factor V. *Proc. Natl. Acad. Sci. USA* 84: 4846-4850.
4. Davie, E.W., Fujikawa, K. and Kisiel, W. 1991. The coagulation cascade: initiation, maintenance, and regulation. *Biochemistry* 30: 10363-10370.
5. Rand, M.D., Kalafatis, M. and Mann, K.G. 1994. Platelet coagulation Factor Va: the major secretory platelet phosphoprotein. *Blood* 83: 2180-2190.
6. Macedo-Ribeiro, S., Bode, W., Huber, R., Quinn-Allen, M.A., Kim, S.W., Ortel, T.L., Bourenkov, G.P., Bartunik, H.D., Stubbs, M.T., Kane, W.H. and Fuentes-Prior, P. 1999. Crystal structures of the membrane-binding C2 domain of human coagulation Factor V. *Nature* 402: 434-439.
7. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 227400. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: F5 (human) mapping to 1q24.2.

PRODUCT

Factor V (h3): 293T Lysate represents a lysate of human Factor V transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Factor V (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive Factor V antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.