

Osteoglycin (h): 293 Lysate: sc-129364

BACKGROUND

The small leucine-rich proteoglycan (SLRP) family of proteins contains various proteins such as Decorin, Biglycan, Fibromodulin, Keratocan, Lumican, Osteoadherin and Osteoglycin. These proteins all have similar functions as they all mediate extracellular matrix organization and act as binding partners of TGF β . Osteoglycin, which also may be designated osteoinductive factor (OIF), is a secreted protein detected in bone tissues. Osteoglycin induces the formation of bone in conjunction with either TGF β 1 or TGF β 2. The precursor form of the OGN gene product, designated Mimecan, is subject to *in situ* proteolytic cleavage to yield the mature Osteoglycin.

REFERENCES

1. Tasheva, E.S., Corpuz, L.M., Funderburgh, J.L. and Conrad, G.W. 1997. Differential splicing and alternative polyadenylation generate multiple Mimecan mRNA transcripts. *J. Biol. Chem.* 272: 32551-32556.
2. Tasheva, E.S., Funderburgh, M.L., McReynolds, J., Funderburgh, J.L. and Conrad, G.W. 1999. The bovine Mimecan gene. Molecular cloning and characterization of two major RNA transcripts generated by alternative use of two splice acceptor sites in the third exon. *J. Biol. Chem.* 274: 18693-18701.
3. Pellegata, N.S., Dieguez-Lucena, J.L., Joensuu, T., Lau, S., Montgomery, K.T., Krahe, R., Kivela, T., Kucherlapati, R., Forsius, H. and de la Chapelle, A. 2000. Mutations in KERA, encoding Keratocan, cause cornea plana. *Nat. Genet.* 25: 91-95.
4. Tasheva, E.S., Klocke, B. and Conrad, G.W. 2004. Analysis of transcriptional regulation of the small leucine rich proteoglycans. *Mol. Vis.* 10: 758-772.
5. Moali, C., Font, B., Ruggiero, F., Eichenberger, D., Rousselle, P., Francois, V., Oldberg, A., Bruckner-Tuderman, L. and Hulmes, D.J. 2005. Substrate-specific modulation of a multi-substrate proteinase. C-terminal processing of fibrillar procollagens is the only BMP-1 dependent activity to be enhanced by PCPE-1. *J. Biol. Chem.* 280: 24188-24194.
6. Wang, X., Ford, B.C., Praul, C.A. and Leach, R.M., Jr. 2005. Characterization of the non-collagenous proteins in avian cortical and medullary bone. *Comp. Biochem. Physiol. B, Biochem. Mol. Biol.* 140: 665-672.
7. Peterson, B., Zhang, J., Iglesias, R., Kabo, M., Hedrick, M., Benhaim, P. and Lieberman, J.R. 2005. Healing of critically sized femoral defects, using genetically modified mesenchymal stem cells from human adipose tissue. *Tissue Eng.* 11: 120-129.
8. Xie, J., Baumann, M.J. and McCabe, L.R. 2005. Adsorption of serum fetuin to hydroxylapatite does not contribute to osteoblast phenotype modifications. *J. Biomed. Mater. Res. A* 73A: 39-47.

CHROMOSOMAL LOCATION

Genetic locus: OGN (human) mapping to 9q22.31.

PRODUCT

Osteoglycin (h): 293 Lysate represents a lysate of human Osteoglycin transfected 293 cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Osteoglycin (h): 293 Lysate is suitable as a Western Blotting positive control for human reactive Osteoglycin antibodies. Recommended use: 10-20 μ l per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.