



Pol II RPB6 (h2): 293T Lysate: sc-129508

BACKGROUND

RNA polymerase II (Pol II) is a multi-subunit enzyme responsible for the transcription of protein-coding genes. Transcription initiation requires recruitment of the complete transcription machinery to a promoter via solicitation by activators and chromatin remodeling factors. Pol II can coordinate 10 to 14 subunits. This complex interacts with the promoter regions of genes and a variety of elements and transcription factors. The DNA binding domain of the polymerase is a groove where TFIIB orients the DNA for unwinding and transcription.

REFERENCES

1. Bushnell, D.A., et al. 2004. Structural basis of transcription: an RNA polymerase II-TFIIB cocrystal at 4.5 Angstroms. *Science* 303: 983-988.
2. Palangat, M., et al. 2004. Downstream DNA selectively affects a paused conformation of human RNA polymerase II. *J. Mol. Biol.* 341: 429-442.
3. Zhong, S., et al. 2004. Epidermal growth factor enhances cellular TATA binding protein levels and induces RNA polymerase I- and III-dependent gene activity. *Mol. Cell. Biol.* 24: 5119-5129.
4. Hirsch, H.A., et al. 2004. Distinct mechanisms for repression of RNA polymerase III transcription by the retinoblastoma tumor suppressor protein. *Mol. Cell. Biol.* 24: 5989-5999.
5. White, R.J. 2004. RNA polymerase III transcription and cancer. *Oncogene* 23: 3208-3216.
6. Cabart, P., et al. 2004. BRCA1 cooperates with NUFIP and P-TEFb to activate transcription by RNA polymerase II. *Oncogene* 23: 5316-5329.
7. Svejstrup, J.Q. 2004. The RNA polymerase II transcription cycle: cycling through chromatin. *Biochim. Biophys. Acta* 1677: 64-73.
8. Cramer, P. 2004. Structure and function of RNA polymerase II. *Adv. Protein Chem.* 67: 1-42.
9. Comai, L. 2004. Mechanism of RNA polymerase I transcription. *Adv. Protein Chem.* 67: 123-155.

CHROMOSOMAL LOCATION

Genetic locus: POLR2F (human) mapping to 22q13.1.

PRODUCT

Pol II RPB6 (h2): 293T Lysate represents a lysate of human Pol II RPB6 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Pol II RPB6 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Pol II RPB6 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.