

# GABP- $\beta$ 1/2 (11): sc-130315

## BACKGROUND

The transcription factor GA-binding protein (GABP) is composed of two subunits, the Ets-related GABP- $\alpha$  and a GABP- $\alpha$ -associated subunit, GABP- $\beta$ . GABP- $\alpha$  binds to a specific DNA sequence and GABP- $\beta$  exists as  $\beta$ 1 and  $\beta$ 2 splice variants that differ in their C-termini. In primary neuronal cultures, GABP- $\beta$  is expressed in both the cytoplasm and the nucleus, whereas GABP- $\alpha$  is expressed mainly in the nucleus. GABP is constitutively expressed as either a GABP- $\alpha\beta$  heterodimer or a GABP- $\alpha\beta$  heterotetramer, both of which can modify GABP-dependent transcription *in vitro* and *in vivo*. The GABP- $\alpha\beta$  tetrameric complex performs many different functions, such as stimulating transcription of the adenovirus E4 gene, differentially activating BRCA1 expression in human breast cell lines, potentiating Tat-mediated activation of long terminal repeat promoter transcription and viral replication in certain cell types, acting as a coordinator of mitochondrial and nuclear transcription for cytochrome oxidase in neurons and assisting in the regulation of rpl32 gene transcription.

## REFERENCES

- Martin, M.E., et al. 1996. Redox regulation of GA-binding protein- $\alpha$  DNA binding activity. *J. Biol. Chem.* 271: 25617-25623.
- Suzuki, F., et al. 1998. Functional interactions of transcription factor human GA-binding protein subunits. *J. Biol. Chem.* 273: 29302-29308.
- Sawada, J., et al. 1999. Synergistic transcriptional activation by hGABP and select members of the activation transcription factor/cAMP response element-binding protein family. *J. Biol. Chem.* 274: 35475-35482.
- Hoare, S., et al. 1999. Identification of a GABP- $\alpha/\beta$  binding site involved in the induction of oxytocin receptor gene expression in human breast cells, potentiation by c-Fos/c-Jun. *Endocrinology* 140: 2268-2279.
- Verhoef, K., et al. 1999. Evolution of the human immunodeficiency virus type 1 long terminal repeat promoter by conversion of an NF $\kappa$ B enhancer element into a GABP binding site. *J. Virol.* 73: 1331-1340.
- Atlas, E., et al. 2000. GA-binding protein  $\alpha/\beta$  is critical regulator of the BRCA1 promoter. *Oncogene* 19: 1933-1940.
- Chinenov, Y., et al. 2000. The  $\alpha$  and  $\beta$  subunits of the GA-binding protein form a stable heterodimer in solution. Revised model of heterotetrameric complex assembly. *J. Biol. Chem.* 275: 7749-7756.

## CHROMOSOMAL LOCATION

Genetic locus: GABPB1 (human) mapping to 15q21.2, GABPB2 (human) mapping to 1q21.3; Gabpb1 (mouse) mapping to 2 F1, Gabpb2 (mouse) mapping to 3 F2.1.

## SOURCE

GABP- $\beta$ 1/2 (11) is a mouse monoclonal antibody raised against recombinant GABP- $\beta$ 1/2 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

GABP- $\beta$ 1/2 (11) is recommended for detection of GABP- $\beta$ 1 and GABP- $\beta$ 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GABP- $\beta$ 1/2 siRNA (h): sc-37903, GABP- $\beta$ 1/2 siRNA (m): sc-37904, GABP- $\beta$ 1/2 shRNA Plasmid (h): sc-37903-SH, GABP- $\beta$ 1/2 shRNA Plasmid (m): sc-37904-SH, GABP- $\beta$ 1/2 shRNA (h) Lentiviral Particles: sc-37903-V and GABP- $\beta$ 1/2 shRNA (m) Lentiviral Particles: sc-37904-V.


Molecular Weight of GABP- $\beta$ 1/2: 42 kDa.

Positive Controls: GABP- $\beta$ 1/2 (h): 293T Lysate: sc-113433, Jurkat whole cell lysate: sc-2204 or C6 whole cell lysate: sc-364373.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotting A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



GABP- $\beta$ 1/2 (11): sc-130315. Western blot analysis of GABP- $\beta$ 1/2 expression in Jurkat (A), 3T3-L1 (B), PC-12 (C) and C6 (D) whole cell lysates and HeLa nuclear extract (E).



GABP- $\beta$ 1/2 (11): sc-130315. Western blot analysis of GABP- $\beta$ 1/2 expression in non-transfected: sc-117752 (A) and human GABP- $\beta$ 1/2 transfected: sc-113433 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

- Ripperger, T., et al. 2015. The heteromeric transcription factor GABP activates the ITGAM/CD11b promoter and induces myeloid differentiation. *Biochim. Biophys. Acta* 1849: 1145-1154.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.