

GAD-65 (61/8F9): sc-130569

BACKGROUND

There are two forms of glutamic acid decarboxylases (GADs) that are found in the brain: GAD-65 (also known as GAD2) and GAD-67 (also known as GAD1, GAD or SCP). GAD-65 and GAD-67 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate limiting step in the production of GABA (γ -aminobutyric acid) from L-glutamic acid. Although both GADs are found in the brain, GAD-65 localizes to synaptic vesicle membranes in nerve terminals, while GAD-67 is distributed throughout the cell. GAD-67 is responsible for the basal levels of GABA synthesis. In the case of a heightened demand for GABA in neurotransmission, GAD-65 will transiently activate to assist in GABA production. The loss of GAD-65 is detrimental and can impair GABA neurotransmission, however the loss of GAD-67 is lethal. Due to alternative splicing, two isoforms exist for GAD-67, the predominant GAD-67 form and the minor GAD-25 form. GAD-25 is not expressed in brain but can be found in a variety of endocrine tissues.

REFERENCES

1. Chessler, S.D., et al. 2002. Immune reactivity to GAD-25 in type 1 diabetes mellitus. *Autoimmunity* 35: 335-341.
2. Kanter, I.C., et al. 2007. Cyclophosphamide for anti-GAD antibody-positive refractory status epilepticus. *Epilepsia* 49: 914-920.
3. Korpershoek, E., et al. 2007. Expression of GAD-67 and novel GAD-67 splice variants during human fetal pancreas development: GAD-67 expression in the fetal pancreas. *Endocr. Pathol.* 18: 31-36.
4. Kanaani, J., et al. 2008. A palmitoylation cycle dynamically regulates partitioning of the GABA-synthesizing enzyme GAD-65 between ER-Golgi and post-Golgi membranes. *J. Cell Sci.* 121: 437-449.
5. Wei, J. and Wu, J.Y. 2008. Post-translational regulation of L-glutamic acid decarboxylase in the brain. *Neurochem. Res.* 33: 1459-1465.
6. Hwang, I.K., et al. 2008. Comparison of glutamic acid decarboxylase 67 immunoreactive neurons in the hippocampal CA1 region at various age stages in dogs. *Neurosci. Lett.* 431: 251-255.
7. Ito, T., et al. 2008. Some γ -motoneurons contain γ -aminobutyric acid in the rat cervical spinal cord. *Brain Res.* 1201: 78-87.
8. Hamilton, K.A., et al. 2008. Sensory deafferentation transsynaptically alters neuronal GluR-1 expression in the external plexiform layer of the adult mouse main olfactory bulb. *Chem. Senses* 33: 201-210.
9. Jain, R., et al. 2008. Innocuous IFN- γ induced by adjuvant-free antigen restores normoglycemia in NOD mice through inhibition of IL-17 production. *J. Exp. Med.* 205: 207-218.

CHROMOSOMAL LOCATION

Genetic locus: GAD2 (human) mapping to 10p12.1; Gad2 (mouse) mapping to 2 A3.

SOURCE

GAD-65 (61/8F9) is a mouse monoclonal antibody raised against full-length recombinant GAD-65 of human origin, with epitope mapping to amino acids 4-17.

PRODUCT

Each vial contains 200 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

GAD-65 (61/8F9) is recommended for detection of GAD-65 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500); non cross-reactive with GAD-67.

Suitable for use as control antibody for GAD-65 siRNA (h): sc-41964, GAD-65 siRNA (m): sc-41965, GAD-65 siRNA (r): sc-61888, GAD-65 shRNA Plasmid (h): sc-41964-SH, GAD-65 shRNA Plasmid (m): sc-41965-SH, GAD-65 shRNA Plasmid (r): sc-61888-SH, GAD-65 shRNA (h) Lentiviral Particles: sc-41964-V, GAD-65 shRNA (m) Lentiviral Particles: sc-41965-V and GAD-65 shRNA (r) Lentiviral Particles: sc-61888-V.

Molecular Weight of GAD-65: 65 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **GAD-65 (A-3): sc-377145** for GAD-65 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647.