

TFIIE- α (F-2): sc-133065

BACKGROUND

In eukaryotic systems, initiation of transcription from protein-coding genes is a complex process requiring RNA polymerase II and broad families of auxiliary transcription factors. Such factors can be divided into two major functional classes: the basal factors that are required for transcription of all Pol II genes, including TFIIA, TFIIB, TFIID, TFIIE, TFIIIF and TFIIH; and sequence-specific factors that regulate gene expression. The basal transcription factors and Pol II form a specific multiprotein complex near the transcription start site by interacting with core promoter elements such as the TATA box generally located 25-30 base pairs upstream of the transcription start site. Human TFIIE consists of two subunits, α and β . The structure of TFIIE appears to be a heterotetramer ($\alpha_2\beta_2$); both subunits are required for optimal basal-level transcription.

CHROMOSOMAL LOCATION

Genetic locus: GTF2E1 (human) mapping to 3q13.33; Gtf2e1 (mouse) mapping to 16 B3.

SOURCE

TFIIE- α (F-2) is a mouse monoclonal antibody raised against amino acids 140-439 mapping at the C-terminus of TFIIE- α of human origin.

PRODUCT

Each vial contains 200 μ g IgG κ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-133065 X, 200 μ g/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TFIIE- α (F-2) is recommended for detection of TFIIE- α of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TFIIE- α siRNA (h): sc-36651, TFIIE- α siRNA (m): sc-36652, TFIIE- α shRNA Plasmid (h): sc-36651-SH, TFIIE- α shRNA Plasmid (m): sc-36652-SH, TFIIE- α shRNA (h) Lentiviral Particles: sc-36651-V and TFIIE- α shRNA (m) Lentiviral Particles: sc-36652-V.

TFIIE- α (F-2) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TFIIE- α : 57 kDa.

Positive Controls: NIH/3T3 nuclear extract: sc-2138, 3611-RF nuclear extract: sc-2143 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



TFIIE- α (F-2): sc-133065. Western blot analysis of TFIIE- α expression in HeLa (A) and A-431 (B) whole cell lysates and Jurkat (C), 3611-RF (D) and NIH/3T3 (E) nuclear extracts.

SELECT PRODUCT CITATIONS

- Gosmain, Y., et al. 2010. Pax6 controls the expression of critical genes involved in pancreatic α cell differentiation and function. *J. Biol. Chem.* 285: 33381-33393.
- Bunch, H., et al. 2021. BRCA1-BARD1 regulates transcription through modulating topoisomerase II β . *Open Biol.* 11: 210221.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.