

SEN3 (G-3): sc-133149

BACKGROUND

SUMO (small ubiquitin-related modifier), a member of the ubiquitin-like protein family, regulates diverse cellular functions of a variety of target proteins, including transcription, DNA repair, nucleocytoplasmic trafficking and chromosome segregation. SUMO precursor proteins undergo cleavage of the residues after the "GG" region by SUMO-specific proteases in maturation. This cleavage of the precursor is a prerequisite for subsequent sumoylation. The sentrin-specific (or SUMO-specific) protease (SEN3) proteins belong to the peptidase C48 family and include SEN1-3 and SEN5-8. SEN1, SEN2 and SEN3 degrade UBL1 and SMT3H2 conjugates and subsequently release the monomers from sumoylated substrates. HIPK2 is a desumoylation target for SEN1 which shuttles between the cytoplasm and the nucleus. Mutation analyses reveal that SEN1 contains the nuclear export sequence (NES) within the extreme carboxyl-terminal region, and SEN1 is exported to the cytoplasm in a NES-dependent manner. SEN2 has been implicated as a downregulator of CTNNB1 levels and may therefore be a modulator of the Wnt pathway. SUMO protease SEN3 reverses the sumoylation of MEF-2 to augment its transcriptional and myogenic activities. SEN5 localizes to the nucleolus and preferentially processes SUMO-3. It is thought to play a role in mitosis and/or cytokinesis. SEN6 localizes to the cytoplasm and releases SUMO-1. Expression of SEN6 is higher in reproductive organs, indicating that it may mediate processes related to reproduction. SEN8 is involved in the release of sentrins.

CHROMOSOMAL LOCATION

Genetic locus: SEN3 (human) mapping to 17p13.1; Senp3 (mouse) mapping to 11 B3.

SOURCE

SEN3 (G-3) is a mouse monoclonal antibody raised against acids 195-389 mapping within an internal region of SEN3 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

SEN3 (G-3) is recommended for detection of SEN3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SEN3 siRNA (h): sc-44451, SEN3 siRNA (m): sc-45718, SEN3 shRNA Plasmid (h): sc-44451-SH, SEN3 shRNA Plasmid (m): sc-45718-SH, SEN3 shRNA (h) Lentiviral Particles: sc-44451-V and SEN3 shRNA (m) Lentiviral Particles: sc-45718-V.


Molecular Weight of SEN3: 72 kDa.

Positive Controls: SEN3 (h): 293T Lysate: sc-111446, HeLa nuclear extract: sc-2120 or PC-3 nuclear extract: sc-2152.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



SEN3 (G-3): sc-133149. Western blot analysis of SEN3 expression in non-transfected: sc-117752 (A) and human SEN3 transfected: sc-111446 (B) 293T whole cell lysates.



SEN3 (G-3): sc-133149. Immunofluorescence staining of formalin-fixed HepG2 cells showing nuclear localization.

SELECT PRODUCT CITATIONS

- Finkbeiner, E., et al. 2011. The SUMO system controls nucleolar partitioning of a novel mammalian ribosome biogenesis complex. *EMBO J.* 30: 1067-1078.
- Wang, Y., et al. 2012. The biphasic redox sensing of SEN3 accounts for the HIF-1 transcriptional activity shift by oxidative stress. *Acta Pharmacol. Sin.* 33: 953-963.
- Jin, S., et al. 2022. Suppression of ACE2 SUMOylation protects against SARS-CoV-2 infection through TOLLIP-mediated selective autophagy. *Nat. Commun.* 13: 5204.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.