

# apoC-IV (7-RE36): sc-134263

## BACKGROUND

Apolipoproteins are protein components of plasma lipoproteins. The apolipoprotein C gene family encodes four homologous proteins designated apoC-I to -IV, which specifically modulate the metabolism of triglyceride-rich lipoproteins. The human apoC-I gene maps to chromosome 19q13.2 and is expressed primarily in the liver, where it is activated when monocytes differentiate into macrophages. The human apoC-II gene maps to chromosome 19q13.2 and encodes a 79 amino acid single chain protein that is a necessary cofactor for the activation of lipoprotein lipase, the enzyme that hydrolyzes triglycerides in plasma and transfers the fatty acids to tissues. The human apoC-III gene maps to chromosome 11q23 and encodes a protein that may delay catabolism of triglyceride-rich particles by inhibiting lipoprotein lipase and hepatic lipase. The human apoC-IV gene maps to chromosome 19q13.32 and encodes a 127 amino acid protein that is primarily expressed in the liver.

## REFERENCES

- Breckenridge, W.C., Little, J.A., Steiner, G., Chow, A. and Poapst, M. 1978. Hypertriglyceridemia associated with deficiency of apolipoprotein C-II. *N. Engl. J. Med.* 298: 1265-1273.
- Allan, C.M., Walker, D., Segrest, J.P. and Taylor, J.M. 1995. Identification and characterization of a new human gene (apoC4) in the apolipoprotein E, C-I and C-II gene locus. *Genomics* 28: 291-300.
- Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 207750. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Jong, M.C. and Havekes, L.M. 2000. Insights into apolipoprotein C metabolism from transgenic and gene-targeted mice. *Int. J. Tissue React.* 22: 59-66.
- Mak, P.A., Laffitte, B.A., Desrumaux, C., Joseph, S.B., Curtiss, L.K., Mangelsdorf, D.J., Tontonoz, P. and Edwards, P.A. 2002. Regulated expression of the apolipoprotein E/C-I/C-IV/C-II gene cluster in murine and human macrophages. A critical role for nuclear liver X receptors  $\alpha$  and  $\beta$ . *J. Biol. Chem.* 277: 31900-31908.
- Kotite, L., Zhang, L.H., Yu, Z., Burlingame, A.L. and Havel, R.J. 2003. Human apoC-IV: isolation, characterization and immunochemical quantification in plasma and plasma lipoproteins. *J. Lipid Res.* 44: 1387-1394.

## CHROMOSOMAL LOCATION

Genetic locus: APOC4 (human) mapping to 19q13.32.

## SOURCE

apoC-IV (7-RE36) is a mouse monoclonal antibody raised against recombinant apoC-IV protein of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

apoC-IV (7-RE36) is recommended for detection of apoC-IV of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for apoC-IV siRNA (h): sc-41187, apoC-IV shRNA Plasmid (h): sc-41187-SH and apoC-IV shRNA (h) Lentiviral Particles: sc-41187-V.


Molecular Weight of apoC-IV: 16 kDa.

Positive Controls: human apoC-IV transfected 293T whole cell lysate.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



apoC-IV (7-RE36): sc-134263. Western blot analysis of apoC-IV expression in human apoC-IV transfected (A) and non-transfected (B) 293T whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.