

USP2 (H-50): sc-135282

BACKGROUND

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove Ub, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP2 (Ub specific peptidase 2), also known as 41 kDa Ub-specific protease, is a 605 amino acid deubiquitinating enzyme that participates in the Ub pathway. Localized to the cytoplasm, USP2 forms a homooligomer and catalyzes the reaction of the Ub C-terminal thioester with water to form Ub and a thiol. USP2 is expressed as two isoforms produced by alternative splicing.

REFERENCES

- Gewies, A. and Grimm, S. 2003. UBP41 is a proapoptotic ubiquitin-specific protease. *Cancer Res.* 63: 682-688.
- Graner, E., Tang, D., Rossi, S., Baron, A., Migita, T., Weinstein, L.J., Lechpammer, M., Huesken, D., Zimmermann, J., Signoretti, S. and Loda, M. 2004. The isopeptidase USP2a regulates the stability of fatty acid synthase in prostate cancer. *Cancer Cell* 5: 253-261.
- Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 604725. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Renatus, M., Parrado, S.G., D'Arcy, A., Eidhoff, U., Gerhartz, B., Hassiepen, U., Pierrat, B., Riedl, R., Vinzenz, D., Worpenberg, S. and Kroemer, M. 2006. Structural basis of ubiquitin recognition by the deubiquitinating protease USP2. *Structure* 14: 1293-1302.
- Horton, R.A., Strachan, E.A., Vogel, K.W. and Riddle, S.M. 2007. A substrate for deubiquitinating enzymes based on time-resolved fluorescence resonance energy transfer between terbium and yellow fluorescent protein. *Anal. Biochem.* 360: 138-143.
- Stevenson, L.F., Sparks, A., Allende-Vega, N., Xirodimas, D.P., Lane, D.P. and Saville, M.K. 2007. The deubiquitinating enzyme USP2a regulates the p53 pathway by targeting MDM2. *EMBO J.* 26: 976-986.

CHROMOSOMAL LOCATION

Genetic locus: USP2 (human) mapping to 11q23.3; Usp2 (mouse) mapping to 9 A5.1.

SOURCE

USP2 (H-50) is a rabbit polyclonal antibody raised against amino acids 501-550 mapping within an internal region of USP2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

USP2 (H-50) is recommended for detection of USP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

USP2 (H-50) is also recommended for detection of USP2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for USP2 siRNA (h): sc-76821, USP2 siRNA (m): sc-76822, USP2 shRNA Plasmid (h): sc-76821-SH, USP2 shRNA Plasmid (m): sc-76822-SH, USP2 shRNA (h) Lentiviral Particles: sc-76821-V and USP2 shRNA (m) Lentiviral Particles: sc-76822-V.


Molecular Weight of USP2: 68 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, HeLa whole cell lysate: sc-2200 or HCT-116 whole cell lysate: sc-364175.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



USP2 (H-50): sc-135282. Western blot analysis of USP2 expression in HL-60 (A), HeLa (B), K-562 (C), HCT-116 (D) and SK-BR-3 (E) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.