

# Ubr2 (8H10): sc-135594

## BACKGROUND

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). Ubr2 (ubiquitin-protein ligase E3- $\alpha$ -2), also known as N-recogin-2, is a 1,755 amino acid protein that contains one UBR-type zinc finger and one RING-type zinc finger. Participating in protein modification events within the N-end rule pathway, Ubr2 functions as an E3 ubiquitin-protein ligase that recognizes and binds proteins that contain destabilizing N-terminal residues, thereby leading to their ubiquitination and subsequent degradation. Mice lacking Ubr2 are infertile due to defects in male meiosis.

## REFERENCES

1. Tasaki, T., et al. 2005. A family of mammalian E3 ubiquitin ligases that contain the UBR box motif and recognize N-degrons. *Mol. Cell. Biol.* 25: 7120-7136.
2. Ouyang, Y., et al. 2006. Loss of Ubr2, an E3 ubiquitin ligase, leads to chromosome fragility and impaired homologous recombinational repair. *Mutat. Res.* 596: 64-75.
3. An, J.Y., et al. 2006. Impaired neurogenesis and cardiovascular development in mice lacking the E3 ubiquitin ligases Ubr1 and Ubr2 of the N-end rule pathway. *Proc. Natl. Acad. Sci. USA* 103: 6212-6217.
4. Lee, M.J., et al. 2008. Synthetic heterovalent inhibitors targeting recognition E3 components of the N-end rule pathway. *Proc. Natl. Acad. Sci. USA* 105: 100-105.
5. Tasaki, T., et al. 2009. The substrate recognition domains of the N-end rule pathway. *J. Biol. Chem.* 284: 1884-1895.
6. Kume, K., et al. 2010. Role of N-end rule ubiquitin ligases Ubr1 and Ubr2 in regulating the leucine-mTOR signaling pathway. *Genes Cells* 15: 339-349.
7. An, J.Y., et al. 2010. UBR2 mediates transcriptional silencing during spermatogenesis via histone ubiquitination. *Proc. Natl. Acad. Sci. USA* 107: 1912-1917.

## CHROMOSOMAL LOCATION

Genetic locus: UBR2 (human) mapping to 6p21.1.

## SOURCE

Ubr2 (8H10) is a mouse monoclonal antibody raised against recombinant Ubr2 protein of human origin.

## PRODUCT

Each vial contains 50  $\mu$ g IgG<sub>2a</sub> kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Ubr2 (8H10) is recommended for detection of Ubr2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Ubr2 siRNA (h): sc-95444, Ubr2 shRNA Plasmid (h): sc-95444-SH and Ubr2 shRNA (h) Lentiviral Particles: sc-95444-V.


Molecular Weight of Ubr2: 201/50/66 kDa.

Positive Controls: human Ubr2 transfected 293T whole cell lysate.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



Ubr2 (8H10): sc-135594. Western blot analysis of Ubr2 expression in human Ubr2 transfected (A) and non-transfected (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

1. Eldeeb, M.A. and Fahlman, R.P. 2014. The anti-apoptotic form of tyrosine kinase Lyn that is generated by proteolysis is degraded by the N-end rule pathway. *Oncotarget* 5: 2714-2722.
2. Eldeeb, M.A. and Fahlman, R.P. 2016. Phosphorylation impacts N-end rule degradation of the proteolytically activated form of BMX kinase. *J. Biol. Chem.* 291: 22757-22768.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.