

G_{αs} (12): sc-135914

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (e.g., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein α , β and γ polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their α subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Evidence, however, has established an important regulatory role for the $\beta\gamma$ subunits. The G_s sub-family of G_α subunits includes two closely related proteins, G_{αs} and G_{αolf}, which respectively stimulate adenylyl cyclase and mediate response to olfactory stimuli.

CHROMOSOMAL LOCATION

Genetic locus: GNAS (human) mapping to 20q13.32; Gnas (mouse) mapping to 2 H4.

SOURCE

G_{αs} (12) is a mouse monoclonal antibody raised against amino acids 11-21 of G_{αs} of human origin.

PRODUCT

Each vial contains 200 μg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

G_{αs} (12) is available conjugated to agarose (sc-135914 AC), 500 μg/0.25 ml agarose in 1 ml, for IP; and to HRP (sc-135914 HRP), 200 μg/ml, for WB, IHC(P) and ELISA.

APPLICATIONS

G_{αs} (12) is recommended for detection of G_{αs} of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for G_{αs} siRNA (h): sc-29328, G_{αs} siRNA (m): sc-41757, G_{αs} shRNA Plasmid (h): sc-29328-SH, G_{αs} shRNA Plasmid (m): sc-41757-SH, G_{αs} shRNA (h) Lentiviral Particles: sc-29328-V and G_{αs} shRNA (m) Lentiviral Particles: sc-41757-V.

Molecular Weight of G_{αs}: 49 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, TT whole cell lysate: sc-364195 or 3T3-L1 cell lysate: sc-2243.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



G_{αs} (12): sc-135914. Western blot analysis of G_{αs} expression in 3T3-L1 (A), C3H/10T1/2 (B) and C6 (C) whole cell lysates and rat brain (D), rat cerebellum (E) and mouse brain (F) tissue extracts.



G_{αs} (12): sc-135914. Western blot analysis of G_{αs} expression in Jurkat (A), HEK293 (B), TT (C), K-562 (D) and MCF7 (E) whole cell lysates.

SELECT PRODUCT CITATIONS

- Lee, I.T., et al. 2012. Role of TLR4/NADPH oxidase/ROS-activated p38 MAPK in VCAM-1 expression induced by lipopolysaccharide in human renal mesangial cells. *Cell Commun. Signal.* 10: 33.
- Tobar-Rubin, R., et al. 2013. Intragenic suppression of a constitutively active allele of G_{αs} associated with McCune-Albright syndrome. *J. Mol. Endocrinol.* 50: 193-201.
- Liu, Y., et al. 2017. Dibutylryl-cAMP attenuates pulmonary fibrosis by blocking myofibroblast differentiation via PKA/CREB/CBP signaling in rats with silicosis. *Respir. Res.* 18: 38.
- Pusapati, G.V., et al. 2018. G protein-coupled receptors control the sensitivity of cells to the morphogen Sonic hedgehog. *Sci. Signal.* 11: eaao5749.
- Fish, E.W., et al. 2019. Cannabinoids exacerbate alcohol teratogenesis by a CB1-hedgehog interaction. *Sci. Rep.* 9: 16057.
- Kankanamge, D., et al. 2019. G protein α_q exerts expression level-dependent distinct signaling paradigms. *Cell. Signal.* 58: 34-43.
- Nunez, F.J., et al. 2019. Glucocorticoids rapidly activate cAMP production via G_{αs} to initiate non-genomic signaling that contributes to one-third of their canonical genomic effects. *FASEB J.* 34: 2882-2895.
- Onopiuk, M., et al. 2020. Control of PTH secretion by the TRPC1 ion channel. *JCI Insight* 5: e132496.
- Cheng, C.Y., et al. 2020. Nrf2/HO-1 partially regulates cytoprotective effects of carbon monoxide against urban particulate matter-induced inflammatory responses in oral keratinocytes. *Cytokine* 133: 155185.
- Maziarz, M., et al. 2020. Revealing the activity of trimeric G-proteins in live cells with a versatile biosensor design. *Cell*. E-published.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.