

ICAM-1 (28): sc-135946

BACKGROUND

Cell adhesion molecules (CAMs) are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play important, yet separate, roles in embryogenesis and development. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the $\beta 2\alpha 1$ and $\beta 2\alpha M$ Integrins. ICAM-2 functions as a ligand for lymphocyte function-associated antigen-1 (LFA-1) and is involved in leukocyte adhesion. ICAM-3 is highly expressed on the surface of human eosinophils and, when bound to ligand, may inhibit eosinophil inflammatory responses and survival. ICAM-4, also known as LW glycoprotein, interacts with Integrins $\alpha L\beta 2$, $\alpha M\beta 2$, $\alpha 4\beta 1$, the αV family and $\alpha IIb\beta 3$, and selective binding to different integrins may be relevant to the pathology in a number of red blood cell associated diseases. Lastly, ICAM-5, expressed on telencephalic neurons, binds CD11a/CD18 and thus may act as an adhesion molecule for leukocyte binding in the central nervous system.

REFERENCES

1. Jorgensen, O.S. 1995. Neural cell adhesion molecule (NCAM) as a quantitative marker in synaptic remodeling. *Neurochem. Res.* 20: 533-547.
2. Edelman, G.M. and Jones, F.S. 1995. Developmental control of N-CAM expression by Hox and Pax gene products. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 349: 305-312.
3. Briskin, M.J., et al. 1996. Structural requirements for mucosal vascular addressin binding to its lymphocyte receptor $\alpha 4\beta 7$. Common themes among integrin-Ig family interactions. *J. Immunol.* 156: 719-726.
4. Heiska, L., et al. 1996. Binding of the cytoplasmic domain of intercellular adhesion molecule-2 (ICAM-2) to α -actinin. *J. Biol. Chem.* 271: 26214-26219.
5. Tian, L., et al. 2000. Binding of T lymphocytes to hippocampal neurons through ICAM-5 (telencephalin) and characterization of its interaction with the leukocyte integrin CD11a/CD18. *Eur. J. Immunol.* 30: 810-818.
6. Lu, T.T. and Cyster, J.G. 2002. Integrin-mediated long-term B cell retention in the splenic marginal zone. *Science* 297: 409-412.
7. Zuccarello, D., et al. 2002. Familial chronic nail candidiasis with ICAM-1 deficiency: a new form of chronic mucocutaneous candidiasis. *J. Med. Genet.* 39: 671-675.
8. Kessel, J.M., et al. 2003. Ligation of intercellular adhesion molecule 3 inhibits GM-CSF production by human eosinophils. *J. Allergy Clin. Immunol.* 111: 1024-1031.
9. Mankelov, T.J., et al. 2003. Identification of critical amino-acid residues on the erythroid intercellular adhesion molecule-4 (ICAM-4) mediating adhesion to αV Integrins. *Blood* 103: 1503-1508.

CHROMOSOMAL LOCATION

Genetic locus: ICAM1 (human) mapping to 19p13.2.

SOURCE

ICAM-1 (28) is a mouse monoclonal antibody raised against amino acids 46-160 of ICAM-1 of human origin.

PRODUCT

Each vial contains 50 μ g IgG₁ in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


ICAM-1 (28) is recommended for detection of ICAM-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for ICAM-1 siRNA (h): sc-29354, ICAM-1 shRNA Plasmid (h): sc-29354-SH and ICAM-1 shRNA (h) Lentiviral Particles: sc-29354-V.


Molecular Weight of ICAM-1: 85-110 kDa.

Positive Controls: HCT-8 cell lysate: sc-24675, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.

DATA



ICAM-1 (28): sc-135946. Western blot analysis of ICAM-1 expression in Jurkat (A) and TNF treated U-937 (B) whole cell lysates.



ICAM-1 (28): sc-135946. Western blot analysis of ICAM-1 expression in HCT-8 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **ICAM-1 (G-5): sc-8439** for ICAM-1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.