

# PP2A-A $\alpha$ / $\beta$ (4G7): sc-13600

## BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified, designated PP1, PP2A, PP2B (calcineurin) and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4) is a putative member of a novel PP family. The PP2A family comprises subfamily members PP2A $\alpha$  and PP2A $\beta$ . The PP2A catalytic subunit associates with a variety of regulatory subunits. Regulatory subunits include PP2A-A $\alpha$  and -A $\beta$ , PP2A-B $\alpha$  and -B $\beta$ , PP2A-C $\alpha$  and -C $\beta$  and PP2A-B56 $\alpha$  and -B56 $\beta$ .

## CHROMOSOMAL LOCATION

Genetic locus: PPP2R1A (human) mapping to 19q13.41, PPP2R1B (human) mapping to 11q23.1; Ppp2r1a (mouse) mapping to 17 A3.2, Ppp2r1b (mouse) mapping to 9 A5.3.

## SOURCE

PP2A-A $\alpha$ / $\beta$  (4G7) is a mouse monoclonal antibody raised against a synthetic peptide corresponding to the N-terminal region of human PP2A-A.

## PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PP2A-A $\alpha$ / $\beta$  (4G7) is available conjugated to agarose (sc-13600 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-13600 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-13600 PE), fluorescein (sc-13600 FITC), Alexa Fluor<sup>®</sup> 488 (sc-13600 AF488), Alexa Fluor<sup>®</sup> 546 (sc-13600 AF546), Alexa Fluor<sup>®</sup> 594 (sc-13600 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-13600 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-13600 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-13600 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

PP2A-A $\alpha$ / $\beta$  (4G7) is recommended for detection of PP2A-A $\alpha$  and PP2A-A $\beta$  of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of PP2A-A $\alpha$ : 55 kDa.


Molecular Weight of PP2A-A $\beta$ : 65 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, T-47D cell lysate: sc-2293 or NIH/3T3 whole cell lysate: sc-2210.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



PP2A-A $\alpha$ / $\beta$  (4G7): sc-13600. Western blot analysis of PP2A-A $\alpha$ / $\beta$  expression in T-47D (A), Jurkat (B), P19 (C), NIH/3T3 (D), KNRK (E) and PC-12 (F) whole cell lysates.



PP2A-A $\alpha$ / $\beta$  (4G7): sc-13600. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of exocrine glandular cells and cytoplasmic and nuclear staining of islets of Langerhans (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of exocrine glandular cells and islet cell. Kindly provided by The Swedish Human Protein Atlas (HPA) program (B).

## SELECT PRODUCT CITATIONS

- Baroja, M.L., et al. 2002. Inhibition of CTLA-4 function by the regulatory subunit of serine/threonine phosphatase 2A. *J. Immunol.* 168: 5070-5078.
- Liu, W., et al. 2012. Isoflurane-induced spatial memory impairment by a mechanism independent of Amyloid- $\beta$  levels and Tau protein phosphorylation changes in aged rats. *Neurol. Res.* 34: 3-10.
- Nakashima, A., et al. 2013. A positive role of mammalian Tip41-like protein, TIPRL, in the amino-acid dependent mTORC1-signaling pathway through interaction with PP2A. *FEBS Lett.* 587: 2924-2929.
- Trakhtenberg, E.F., et al. 2014. Regulating set- $\beta$ 's subcellular localization toggles its function between inhibiting and promoting axon growth and regeneration. *J. Neurosci.* 34: 7361-7374.
- Juraszek, B. and Nafecz, K.A. 2016. Protein phosphatase PP2A—a novel interacting partner of carnitine transporter OCTN2 (SLC22A5) in rat astrocytes. *J. Neurochem.* 139: 537-551.
- Göder, A., et al. 2018. HDAC1 and HDAC2 integrate checkpoint kinase phosphorylation and cell fate through the phosphatase-2A subunit PR130. *Nat. Commun.* 9: 764.
- Petersen, C.H., et al. 2019. Possible predisposition for colorectal carcinogenesis due to altered gene expressions in normal appearing mucosa from patients with colorectal neoplasia. *BMC Cancer* 19: 643.
- Kukreti, H. and Amuthavalli, K. 2020. MicroRNA-34a causes ceramide accumulation and effects Insulin signaling pathway by targeting ceramide kinase (CERK) in aging skeletal muscle. *J. Cell. Biochem.* E-published.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.