

SNX1 (51): sc-136247

BACKGROUND

Sorting nexin 1 (SNX1) is a member of a large family of hydrophilic proteins that interact with a variety of receptor types and are involved in intracellular trafficking. SNX1 and the related splice variant, SNX1A, bind the epidermal growth factor (EGF) receptor, facilitate its transport to lysosome, and thereby contribute to the degradation of the receptor. SNX2 and SNX4 share a high degree of amino acid similarity with SNX1, as they all contain a characteristic phox homology (PX) domain. These proteins are all partially associated with cellular membranes, and they, likewise, associate with EGF, PDGF and Insulin receptor tyrosine kinases. These nexins are widely expressed and yet have various tissue distribution patterns. Additionally, the sorting nexins can associate with each other and with a variety of other cellular proteins, suggesting that they exist as part of multisubunit complexes. The related protein, SNX3, comprises a distinct subgroup of nexins that share less sequence similarity outside of the PX domain and have dramatically different binding affinities for the tyrosine kinase receptors.

REFERENCES

1. Trowbridge, I.S., et al. 1993. Signal-dependent membrane protein trafficking in the endocytic pathway. *Annu. Rev. Cell Biol.* 9: 129-161.
2. Opresko, L.K., et al. 1995. Endocytosis and lysosomal targeting of epidermal growth factor receptors are mediated by distinct sequences independent of the tyrosine kinase domain. *J. Biol. Chem.* 270: 4325-4333.
3. Kurten, R.C., et al. 1996. Enhanced degradation of EGF receptors by a sorting nexin, SNX1. *Science* 272: 1008-1010.
4. Ponting, C.P. 1996. Novel domains in NADPH oxidase subunits, sorting nexins, and PtdIns 3-kinases: binding partners of SH3 domains? *Protein Sci.* 5: 2353-2357.
5. Horazdovsky, B.F., et al. 1997. A sorting nexin-1 homologue, Vps5p, forms a complex with Vps17p and is required for recycling the vacuolar protein-sorting receptor. *Mol. Biol. Cell* 8: 1529-1541.
6. Haft, C.R., et al. 1998. Identification of a family of sorting nexin molecules and characterization of their association with receptors. *Mol. Cell. Biol.* 18: 7278-7287.

CHROMOSOMAL LOCATION

Genetic locus: SNX1 (human) mapping to 15q22.31.

SOURCE

SNX1 (51) is a mouse monoclonal antibody raised against amino acids 1-108 of SNX1 of human origin.

PRODUCT

Each vial contains 50 µg IgG₁ in 0.5 ml of PBS with < 0.1% sodium azide, 0.1% gelatin, 20% glycerol, and 0.04% stabilizer protein.

Blocking peptide available for competition studies, sc-136247 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

SNX1 (51) is recommended for detection of SNX1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).


Suitable for use as control antibody for SNX1 siRNA (h): sc-41345, SNX1 shRNA Plasmid (h): sc-41345-SH and SNX1 shRNA (h) Lentiviral Particles: sc-41345-V.

Molecular Weight (predicted) of SNX1: 60 kDa.


Molecular Weight (observed) of SNX1: 78 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

DATA



SNX1 (51): sc-136247. Western blot analysis of SNX1 expression in HeLa whole cell lysate.



SNX1 (51): sc-136247. Immunofluorescence staining of HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

1. Itai, N., et al. 2018. The phosphorylation of sorting nexin 5 at serine 226 regulates retrograde transport and macropinocytosis. *PLoS ONE* 13: e0207205.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures. Not for resale.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.