

Perforin 1 (F-1): sc-136994

BACKGROUND

The body's major defense against virus-infected and tumorigenic cells is cytotoxic T lymphocyte (CTL)-mediated cytotoxicity, which also plays a role in autoimmune diseases and transplant rejection. During CTL-mediated cytotoxicity, CTL granules containing Perforin are exocytosed. Perforin is a pore-forming protein that facilitates the entry of cytotoxic serine proteases, such as granzymes, into target cells by forming transmembrane channels in target cell membranes. Perforin is primarily expressed in cytotoxic T lymphocytes (CTL) and natural killer (NK) cells, but has more recently been observed in an astrocyte population of the human brain. It has been shown that abrogation of Perforin function by Ca^{2+} -complexing agents leads to decreased levels of necrosis, demonstrating that both necrosis and apoptosis contribute to CTL-mediated cytotoxicity. Perforin activity has been shown to be induced by IL-2, IL-3, IL-4, IL-6 and to a lesser degree, TNF and IFN- γ .

REFERENCES

1. Liu, C.C., et al. 1990. Induction of Perforin and serine esterases in a murine cytotoxic T lymphocyte clone. *J. Immunol.* 144: 1196-1201.
2. Podack, E.R., et al. 1991. A central role of Perforin in cytolysis? *Annu. Rev. Immunol.* 9: 129-157.
3. Thia, K.Y., et al. 1993. Expression of human Perforin in a mouse cytotoxic T lymphocyte cell line: evidence for perturbation of granule-mediated cytotoxicity. *J. Leukoc. Biol.* 54: 528-533.

CHROMOSOMAL LOCATION

Genetic locus: PRF1 (human) mapping to 10q22.1; Prf1 (mouse) mapping to 10 B4.

SOURCE

Perforin 1 (F-1) is a mouse monoclonal antibody raised against amino acids 241-555 mapping at the C-terminus of Perforin 1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Perforin 1 (F-1) is available conjugated to agarose (sc-136994 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-136994 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-136994 PE), fluorescein (sc-136994 FITC), Alexa Fluor[®] 488 (sc-136994 AF488), Alexa Fluor[®] 546 (sc-136994 AF546), Alexa Fluor[®] 594 (sc-136994 AF594) or Alexa Fluor[®] 647 (sc-136994 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-136994 AF680) or Alexa Fluor[®] 790 (sc-136994 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Perforin 1 (F-1) is recommended for detection of Perforin 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Perforin 1 siRNA (h): sc-42592, Perforin 1 siRNA (m): sc-42593, Perforin 1 siRNA (r): sc-270073, Perforin 1 shRNA Plasmid (h): sc-42592-SH, Perforin 1 shRNA Plasmid (m): sc-42593-SH, Perforin 1 shRNA Plasmid (r): sc-270073-SH, Perforin 1 shRNA (h) Lentiviral Particles: sc-42592-V, Perforin 1 shRNA (m) Lentiviral Particles: sc-42593-V and Perforin 1 shRNA (r) Lentiviral Particles: sc-270073-V.


Molecular Weight of Perforin 1: 75 kDa.

Positive Controls: Perforin 1 (h): 293T Lysate: sc-116672, CTLL-2 cell lysate: sc-2242 or NK-92 whole cell lysate: sc-364788.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Perforin 1 (F-1): sc-136994. Western blot analysis of Perforin 1 expression in non-transfected 293T: sc-117752 (A), human Perforin 1 transfected 293T: sc-116672 (B), CTLL-2 (C) and NK-92 (D) whole cell lysates.



Perforin 1 (F-1): sc-136994. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse spleen tissue showing cytoplasmic staining of cells in white pulp and cells in red pulp (B).

RESEARCH USE

For research use only, not for use in diagnostic procedures.