

EBF (D-8): sc-137039

BACKGROUND

B lymphocyte maturation is an intricate process that requires a distinct set of transcription factors with respect to the stage of cell differentiation and cell lineage. Among the transcriptional regulators involved in the early stages of B cell development, early B cell factor (EBF), also designated olfactory neuronal transcription factor 1 (OLF1), targets promoter elements for B lymphoid kinase (Blk) and genes encoding portions of the early stage B cell receptors (BCR), which are necessary for initiation of Ig light chain gene recombination and Src kinase (Blk) signaling. EBF is a basic helix-loop-helix (bHLH) homodimeric transcription factor composed of two subunits that interact with the core DNA sequence, CCCNNGGG, through a DNA recognition domain containing a zinc-coordination motif. Promoter elements to certain neuron-specific genes encoding olfactory-related proteins have been shown to contain EBF binding sites.

REFERENCES

1. Wang, M.M., et al. 1993. Molecular cloning of the olfactory neuronal transcription factor OLF1 by genetic selection in yeast. *Nature* 364: 121-126.
2. Lin, H., et al. 1995. Failure of B cell differentiation in mice lacking the transcription factor EBF. *Nature* 376: 263-267.
3. Hagman, J., et al. 1995. EBF contains a novel zinc-coordination motif and multiple dimerization and transcriptional activation domains. *EMBO J.* 14: 2907-2916.
4. Sigvardsson, M., et al. 1997. EBF and E47 collaborate to induce expression of the endogenous immunoglobulin surrogate light chain genes. *Immunity* 7: 25-36.
5. Akerblad, P., et al. 1999. The B29 (immunoglobulin β -chain) gene is a genetic target for early B cell factor. *Mol. Cell. Biol.* 19: 392-401.
6. Akerblad, P., et al. 1999. Early B cell factor is an activator of the B lymphoid kinase promoter in early B cell development. *J. Immunol.* 163: 5453-5461.
7. Sigvardsson, M. 2000. Overlapping expression of early B cell factor and basic helix-loop-helix proteins as a mechanism to dictate B lineage-specific activity of the λ 5 promoter. *Mol. Cell. Biol.* 20: 3640-3654.

SOURCE

EBF (D-8) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of EBF of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

EBF (D-8) is recommended for detection of EBF1, EBF2, EBF3 and EBF4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of EBF: 80 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, A549 cell lysate: sc-2413 or Ramos nuclear extract: sc-2153.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



EBF (D-8): sc-137039. Western blot analysis of EBF expression in A549 (A) and IMR-32 (B) whole cell lysates.



EBF (D-8): sc-137039. Immunoperoxidase staining of formalin fixed, paraffin-embedded human bone marrow tissue showing cytoplasmic staining of hematopoietic cells.

SELECT PRODUCT CITATIONS

1. Rajakumari, S., et al. 2013. EBF2 determines and maintains brown adipocyte identity. *Cell Metab.* 17: 562-574.
2. Dallner, O.S., et al. 2019. Dysregulation of a long noncoding RNA reduces leptin leading to a leptin-responsive form of obesity. *Nat. Med.* 25: 507-516.



See **EBF (C-8): sc-137065** for EBF antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.