

galectin-7 (D-5): sc-137086

BACKGROUND

Galectins are a family of soluble β -galactoside-binding animal lectins that modulate cell-to-cell adhesion and cell-to-extracellular matrix (ECM) interactions and play a role in tumor progression, pre-mRNA splicing and apoptosis. Galectin-7, expressed mainly in stratified squamous epithelium, is activated by p53 and repressed by retinoic acid. It is a pro-apoptotic protein that functions intracellularly upstream of JNK activation and cytochrome c release. The galectin-7 gene maps to chromosome 19.

REFERENCES

1. Couraud, P.O., et al. 1989. Molecular cloning, characterization, and expression of a human 14 kDa lectin. *J. Biol. Chem.* 264: 1310-1316.
2. Hirabayashi, J., et al. 1989. Cloning and nucleotide sequence of a full-length cDNA for human 14 kDa β -galactoside-binding lectin. *Biochim. Biophys. Acta* 1008: 85-91.
3. Madsen, P., et al. 1995. Cloning, expression, and chromosome mapping of human galectin-7. *J. Biol. Chem.* 270: 5823-5829.
4. Magnaldo, T., et al. 1995. Galectin-7, a human 14 kDa S-lectin, specifically expressed in keratinocytes and sensitive to retinoic acid. *Dev. Biol.* 168: 259-271.
5. Magnaldo, T., et al. 1998. Galectin-7, a marker of all types of stratified epithelia. *Differentiation* 63: 159-168.
6. Leonidas, D.D., et al. 1998. Structural basis for the recognition of carbohydrates by human galectin-7. *Biochemistry* 37: 13930-13940.
7. Bernerd, F., et al. 1999. Galectin-7 overexpression is associated with the apoptotic process in UVB-induced sunburn keratinocytes. *Proc. Natl. Acad. Sci. USA* 96: 11329-11334.
8. Kuwabara, I., et al. 2002. Galectin-7 (PIG1) exhibits pro-apoptotic function through JNK activation and mitochondrial cytochrome c release. *J. Biol. Chem.* 277: 3487-3497.
9. Cao, Z., et al. 2002. Galectins-3 and -7, but not galectin-1, play a role in re-epithelialization of wounds. *J. Biol. Chem.* 277: 42299-42305.

CHROMOSOMAL LOCATION

Genetic locus: LGALS7 (human) mapping to 19q13.2.

SOURCE

galectin-7 (D-5) is a mouse monoclonal antibody raised against amino acids 77-136 mapping at the C-terminus of galectin-7 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

galectin-7 (D-5) is recommended for detection of galectin-7 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for galectin-7 siRNA (h): sc-44534, galectin-7 shRNA Plasmid (h): sc-44534-SH and galectin-7 shRNA (h) Lentiviral Particles: sc-44534-V.


Molecular Weight of galectin-7: 14 kDa.

Positive Controls: human cervix extract: sc-363756 or SCC-4 whole cell lysate: sc-364363.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



galectin-7 (D-5): sc-137086. Western blot analysis of galectin-7 expression in human cervix tissue extract (A) and SCC-4 whole cell lysate (B).



galectin-7 (D-5): sc-137086. Immunoperoxidase staining of formalin fixed, paraffin-embedded human oral mucosa tissue showing cytoplasmic staining of squamous epithelial cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.