



# FAM221B siRNA (m): sc-140036

## BACKGROUND

C9orf128 (chromosome 9 open reading frame 128) is a 402 amino acid protein that exists as three alternatively spliced isoforms, which are encoded by a gene that maps to human chromosome 9p13.3. Chromosome 9 consists of about 145 million bases, represents 4% of the human genome and encodes nearly 900 genes. Thought to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of Bcr-Abl fusion protein often found in leukemias.

## REFERENCES

1. Humphray, S.J., Oliver, K., Hunt, A.R., Plumb, R.W., Loveland, J.E., Howe, K.L., Andrews, T.D., Searle, S., Hunt, S.E., Scott, C.E., Jones, M.C., Ainscough, R., Almeida, J.P., Ambrose, K.D., Ashwell, R.I., et al. 2004. DNA sequence and analysis of human chromosome 9. *Nature* 429: 369-374.
2. Coppo, P., Flamant, S., De Mas, V., Jarrier, P., Guillier, M., Bonnet, M.L., Lacout, C., Guilhot, F., Vainchenker, W. and Turhan, A.G. 2006. Bcr-Abl activates Stat3 via JAK and MEK pathways in human cells. *Br. J. Haematol.* 134: 171-179.
3. Zheng, X., Güller, S., Beissert, T., Puccetti, E. and Ruthardt, M. 2006. Bcr and its mutants, the reciprocal t(9;22)-associated Abl/Bcr fusion proteins, differentially regulate the cytoskeleton and cell motility. *BMC Cancer* 7: 262.
4. Burmeister, T., Schwartz, S., Taubald, A., Jost, E., Lipp, T., Schneller, F., Diedrich, H., Thomssen, H., Mey, U.J., Eucker, J., Rieder, H., Gökbuget, N., Hoelzer, D. and Thiel, E. 2007. Atypical Bcr-Abl mRNA transcripts in adult acute lymphoblastic leukemia. *Haematologica* 92: 1699-1702.
5. Cottin, V., Dupuis-Girod, S., Lesca, G. and Cordier, J.F. 2007. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (Rendu-Osler disease). *Respiration* 74: 361-378.
6. Gardiner, J., Barton, D., Marc, J. and Overall, R. 2007. Potential role of Tubulin acetylation and microtubule-based protein trafficking in familial dysautonomia. *Traffic* 8: 1145-1149.
7. Hims, M.M., Shetty, R.S., Pickel, J., Mull, J., Leyne, M., Liu, L., Gusella, J.F. and Slaugenaupt, S.A. 2007. A humanized IKBKAP transgenic mouse models a tissue-specific human splicing defect. *Genomics* 90: 389-396.
8. Temtamy, S.A., Kamel, A.K., Ismail, S., Helmy, N.A., Aglan, M.S., El Gammal, M., El Ruby, M. and Mohamed, A.M. 2007. Phenotypic and cytogenetic spectrum of 9p trisomy. *Genet. Couns.* 18: 29-48.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Fam221b (mouse) mapping to 4 B1.

## PRODUCT

FAM221B siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FAM221B shRNA Plasmid (m): sc-140036-SH and FAM221B shRNA (m) Lentiviral Particles: sc-140036-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

FAM221B siRNA (m) is recommended for the inhibition of FAM221B expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FAM221B gene expression knockdown using RT-PCR Primer: FAM221B (m)-PR: sc-140036-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.