

Unc79 siRNA (m): sc-140499

BACKGROUND

Unc79 (unc-79 homolog (*C. elegans*)) is a 2,635 amino acid multi-pass membrane protein that exists as three alternatively spliced isoforms and is known to interact with NALCN and UNC-80. Unc79 is a component of the NALCN sodium channel complex, which controls neuronal excitability and is activated by neurotensin or neuropeptides substance P. A member of the Unc79 family, Unc79 is encoded by a gene that maps to human chromosome 14q32.13. Chromosome 14 houses over 700 genes and comprises nearly 3.5% of the human genome. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease (AD). The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder α 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction.

REFERENCES

1. Avramopoulos, D., Fallin, M.D. and Bassett, S.S. 2005. Linkage to chromosome 14q in Alzheimer's disease (AD) patients without psychotic symptoms. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 132B: 9-13.
2. Yeh, E., Ng, S., Zhang, M., Bouhours, M., Wang, Y., Wang, M., Hung, W., Aoyagi, K., Melnik-Martinez, K., Li, M., Liu, F., Schafer, W.R. and Zhen, M. 2008. A putative cation channel, NCA-1, and a novel protein, UNC-80, transmit neuronal activity in *C. elegans*. *PLoS Biol.* 6: e55.
3. Lerner, A.J. and Doran, M. 2009. Genotype-phenotype relationships of presenilin-1 mutations in Alzheimer's disease: an update. *J. Alzheimers Dis.* 17: 259-265.
4. Topic, A., Alempijevic, T., Milutinovic, A.S. and Kovacevic, N. 2009. α 1-antitrypsin phenotypes in adult liver disease patients. *Ups. J. Med. Sci.* 114: 228-234.
5. Lu, B., Zhang, Q., Wang, H., Wang, Y., Nakayama, M. and Ren, D. 2010. Extracellular calcium controls background current and neuronal excitability via an UNC79-UNC80-NALCN cation channel complex. *Neuron* 68: 488-499.
6. Specia, D.J., Chihara, D., Ashique, A.M., Bowers, M.S., Pierce-Shimomura, J.T., Lee, J., Rabbee, N., Speed, T.P., Gultarte, R.J., Chitwood, J., Medrano, J.F., Liao, M., Sonner, J.M., Eger, E.I., Peterson, A.S. and McIntire, S.L. 2010. Conserved role of Unc79 in ethanol responses in lightweight mutant mice. *PLoS Genet.* 6 pii: e1001057.
7. Davies, A.G., Friedberg, R.I., Gupta, H., Chan, C.L., Shelton, K.L. and Bettinger, J.C. 2012. Different genes influence toluene- and ethanol-induced locomotor impairment in *C. elegans*. *Drug Alcohol Depend.* 122: 47-54.

CHROMOSOMAL LOCATION

Genetic locus: Unc79 (mouse) mapping to 12 E.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Unc79 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Unc79 shRNA Plasmid (m): sc-140499-SH and Unc79 shRNA (m) Lentiviral Particles: sc-140499-V as alternate gene silencing products.

For independent verification of Unc79 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-140499A and sc-140499B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Unc79 siRNA (m) is recommended for the inhibition of Unc79 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Unc79 gene expression knockdown using RT-PCR Primer: Unc79 (m)-PR: sc-140499-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.