



A530032D15Rik siRNA (m): sc-140652

BACKGROUND

A530032D15Rik (RIKEN cDNA A530032D15 gene), also known as Gm1888 and SP140 nuclear body protein family member, is a 211 amino acid murine protein. SP140 is a component of the nuclear body that may be involved in trafficking between the nucleus and the cytoplasm. SP140 is induced by interferons and contains a bromo domain, a HSR domain, a PHD-type zinc finger and a SAND domain. It is thought that SP140 may participate in the pathogenesis of acute promyelocytic leukemia and viral infection. A530032D15Rik contains one Sp100 domain and is encoded by a gene located on mouse chromosome 1 C5.

REFERENCES

1. Dent, A.L., et al. 1996. LYSP100-associated nuclear domains (LANDs): description of a new class of subnuclear structures and their relationship to PML nuclear bodies. *Blood* 88: 1423-1426.
2. Bloch, D.B., et al. 1996. Identification and characterization of a leukocyte-specific component of the nuclear body. *J. Biol. Chem.* 271: 29198-29204.
3. Bloch, D.B., et al. 1999. Structural and functional heterogeneity of nuclear bodies. *Mol. Cell. Biol.* 19: 4423-4430.
4. Bloch, D.B., et al. 2000. Sp110 localizes to the PML-Sp100 nuclear body and may function as a nuclear hormone receptor transcriptional coactivator. *Mol. Cell. Biol.* 20: 6138-6146.
5. Regad, T., et al. 2001. Role and fate of PML nuclear bodies in response to interferon and viral infections. *Oncogene* 20: 7274-7286.
6. Madani, N., et al. 2002. Implication of the lymphocyte-specific nuclear body protein Sp140 in an innate response to human immunodeficiency virus type 1. *J. Virol.* 76: 11133-11138.
7. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 608602. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Zhou, J.R., et al. 2007. Identification of tumor-associated proteins in laryngeal squamous cell carcinoma by proteomics. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi* 42: 934-938.
9. Granito, A., et al. 2010. PML nuclear body component Sp140 is a novel autoantigen in primary biliary cirrhosis. *Am. J. Gastroenterol.* 105: 125-131.

CHROMOSOMAL LOCATION

Genetic locus: A530032D15Rik (mouse) mapping to 1 C5.

PRODUCT

A530032D15Rik siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see A530032D15Rik shRNA Plasmid (m): sc-140652-SH and A530032D15Rik shRNA (m) Lentiviral Particles: sc-140652-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

A530032D15Rik siRNA (m) is recommended for the inhibition of A530032D15Rik expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor A530032D15Rik gene expression knockdown using RT-PCR Primer: A530032D15Rik (m)-PR: sc-140652-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.