



# Adat3 siRNA (m): sc-140873

## BACKGROUND

Editing of RNA alters the nucleotide sequence of a transcript to produce codon changes, which can result in alternative translation patterns from a single pre-mRNA. One type of RNA editing involves tRNA-specific adenosine deaminase, ADAT1, which is responsible for the first step in the processing of eukaryotic tRNA<sup>Ala</sup> transcripts that undergo specific adenosine to inosine modifications. ADAT2 (tRNA-specific adenosine deaminase 2), also known as deaminase domain-containing protein 1, and ADAT3 (tRNA-specific adenosine deaminase-like protein 3) are also thought to participate in the deamination of adenosine-34 to inosine in many tRNAs. Belonging to the cytidine and deoxycytidylate deaminase protein family, ADAT2 and ADAT3 both employ zinc as a cofactor. ADAT2 is a 191 amino acid protein that exists as two isoforms produced by alternative splicing events. ADAT3 is a 351 amino acid protein that is phosphorylated upon DNA damage, possibly by Atm or ATR.

## REFERENCES

1. Maas, S., et al. 1996. Structural requirements for RNA editing in glutamate receptor pre-mRNAs by recombinant double-stranded RNA adenosine deaminase. *J. Biol. Chem.* 271: 12221-12226.
2. Melcher, T., et al. 1996. RED2, a brain-specific member of the RNA-specific adenosine deaminase family. *J. Biol. Chem.* 271: 31795-31798.
3. Rueter, S.M., et al. 1999. Regulation of alternative splicing by RNA editing. *Nature* 399: 75-80.
4. Maas, S., et al. 1999. Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes. *Proc. Natl. Acad. Sci. USA* 96: 8895-8900.
5. Keller, W., et al. 1999. Editing of messenger RNA precursors and of tRNAs by adenosine to inosine conversion. *FEBS Lett.* 452: 71-76.
6. Schaub, M. and Keller, W. 2002. RNA editing by adenosine deaminases generates RNA and protein diversity. *Biochimie* 84: 791-803.
7. Kimura, K., et al. 2006. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. *Genome Res.* 16: 55-65.
8. Matsuoka, S., et al. 2007. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316: 1160-1166.

## CHROMOSOMAL LOCATION

Genetic locus: Adat3 (mouse) mapping to 10 C1.

## PRODUCT

Adat3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Adat3 shRNA Plasmid (m): sc-140873-SH and Adat3 shRNA (m) Lentiviral Particles: sc-140873-V as alternate gene silencing products.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Adat3 siRNA (m) is recommended for the inhibition of ADAT3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ADAT3 gene expression knockdown using RT-PCR Primer: Adat3 (m)-PR: sc-140873-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.