



# ALG-4 siRNA (m): sc-141008

## BACKGROUND

An increased intracellular  $Ca^{2+}$  concentration induces apoptotic cell death. Transiently elevated  $Ca^{2+}$  concentrations are required for glucocorticoid-mediated and T cell receptor-mediated pathways, leading to T cell apoptosis. ALG-4 (apoptosis-linked gene 4), alternately known as programmed cell death 11 (PDCD11) or NF $\kappa$ B-binding protein (NFBP), is a 1,871 amino acid protein involved in the biosynthesis of rRNA. ALG-4 co-localizes with U3 RNA in the nucleolus and is expressed at moderate levels in testis. ALG-4 is expressed at low levels in lung, liver, placenta, kidney, spleen, skeletal muscle, ovary, peripheral blood leukocytes, colon, small intestine and thymus. ALG-4 contains multiple phosphorylated serine residues, 4 HAT repeats and 12 S<sub>1</sub> motif domains.

## REFERENCES

1. Nagase, T., Seki, N., Ishikawa, K., Tanaka, A. and Nomura, N. 1996. Prediction of the coding sequences of unidentified human genes. V. The coding sequences of 40 new genes (KIAA0161-KIAA0200) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 3: 17-24.
2. Lacana, E. and D'Adamio, L. 1999. Regulation of Fas ligand expression and cell death by apoptosis-linked gene 4. *Nat. Med.* 5: 542-547.
3. Sweet, T., Khalili, K., Sawaya, B.E. and Amini, S. 2003. Identification of a novel protein from glial cells based on its ability to interact with NF $\kappa$ B subunits. *J. Cell. Biochem.* 90: 884-891.
4. Sweet, T., Sawaya, B.E., Khalili, K. and Amini, S. 2005. Interplay between NFBP and NF $\kappa$ B modulates tat activation of the LTR. *J. Cell. Physiol.* 204: 375-380.
5. Sweet, T., Yen, W., Khalili, K. and Amini, S. 2008. Evidence for involvement of NFBP in processing of ribosomal RNA. *J. Cell. Physiol.* 214: 381-388.
6. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612333. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: Pdc11 (mouse) mapping to 19 C3.

## PRODUCT

ALG-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ALG-4 shRNA Plasmid (m): sc-141008-SH and ALG-4 shRNA (m) Lentiviral Particles: sc-141008-V as alternate gene silencing products.

For independent verification of ALG-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141008A, sc-141008B and sc-141008C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ALG-4 siRNA (m) is recommended for the inhibition of ALG-4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ALG-4 gene expression knockdown using RT-PCR Primer: ALG-4 (m)-PR: sc-141008-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.