

AMPD3 siRNA (m): sc-141054

BACKGROUND

Adenosine monophosphate (AMP) deaminase is a cytosolic enzyme responsible for the hydrolytic deamination of AMP to inosine monophosphate (IMP) and NH₃. AMP deaminase functions as a homotetramer and participates in the purine nucleotide cycle, playing an important role in energy metabolism. Three differentially expressed isozymes of AMP deaminase exist in mammals, namely AMPD1, AMPD2 and AMPD3, which is also known as AMP deaminase isoform E or erythrocyte AMP deaminase. The isozymes differ among their N-terminal domains while sharing a conserved C-terminal catalytic domain. AMPD1 is expressed in skeletal muscle; AMPD2 is found in undifferentiated myoblasts, smooth muscle, embryonic muscle and non-muscle tissue; and AMPD3 is expressed in erythrocytes. Encoded by a gene that maps to human chromosome 11p15.4, AMPD3 is a 767 amino acid protein that contains seventeen exons and exists as three alternatively spliced isoforms. AMPD3 mutations can result in the clinically asymptomatic, autosomal recessive condition erythrocyte AMP deaminase deficiency.

REFERENCES

1. Yamada, Y., Goto, H. and Ogasawara, N. 1992. Cloning and nucleotide sequence of the cDNA encoding human erythrocyte-specific AMP deaminase. *Biochim. Biophys. Acta* 1171: 125-128.
2. Mahnke-Zizelman, D.K. and Sabina, R.L. 1992. Cloning of human AMP deaminase isoform E cDNAs. Evidence for a third AMPD gene exhibiting alternatively spliced 5'-exons. *J. Biol. Chem.* 267: 20866-20877.
3. Yamada, Y., Goto, H. and Ogasawara, N. 1994. A point mutation responsible for human erythrocyte AMP deaminase deficiency. *Hum. Mol. Genet.* 3: 331-334.
4. Mahnke-Zizelman, D.K., Eddy, R., Shows, T.B. and Sabina, R.L. 1996. Characterization of the human AMPD3 gene reveals that 5' exon usage is subject to transcriptional control by three tandem promoters and alternative splicing. *Biochim. Biophys. Acta* 1306: 75-92.
5. Mahnke-Zizelman, D.K., D'cunha, J., Wojnar, J.M., Brogley, M.A. and Sabina, R.L. 1997. Regulation of rat AMP deaminase 3 (isoform C) by development and skeletal muscle fibre type. *Biochem. J.* 326: 521-529.
6. Sermsuvitayawong, K., Wang, X., Nagabukuro, A., Matsuda, Y., Morisaki, H., Toyama, K., Mukai, T. and Morisaki, T. 1997. Genomic organization of AMPD3, heart-type AMPD gene, located in mouse chromosome 7. *Mamm. Genome* 8: 767-769.
7. Yamada, Y., Makarewicz, W., Goto, H., Nomura, N., Kitoh, H. and Ogasawara, N. 1998. Gene mutations responsible for human erythrocyte AMP deaminase deficiency in Poles. *Adv. Exp. Med. Biol.* 431: 347-350.
8. Mahnke-Zizelman, D.K., Tullson, P.C. and Sabina, R.L. 1998. Novel aspects of tetramer assembly and N-terminal domain structure and function are revealed by recombinant expression of human AMP deaminase isoforms. *J. Biol. Chem.* 273: 35118-35125.
9. Yamada, Y., Goto, H., Wakamatsu, N. and Ogasawara, N. 2001. A rare case of complete human erythrocyte AMP deaminase deficiency due to two novel missense mutations in AMPD3. *Hum. Mutat.* 17: 78.

CHROMOSOMAL LOCATION

Genetic locus: *Ampd3* (mouse) mapping to 7 F1.

PRODUCT

AMPD3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AMPD3 shRNA Plasmid (m): sc-141054-SH and AMPD3 shRNA (m) Lentiviral Particles: sc-141054-V as alternate gene silencing products.

For independent verification of AMPD3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141054A, sc-141054B and sc-141054C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AMPD3 siRNA (m) is recommended for the inhibition of AMPD3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AMPD3 gene expression knockdown using RT-PCR Primer: AMPD3 (m)-PR: sc-141054-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.