

ArgBP2 siRNA (m): sc-141193

BACKGROUND

Arg and c-Abl represent the mammalian members of the Abelson family of non-receptor protein-tyrosine kinases. They interact with the Arg/Abl binding proteins (ArgBPs) via SH3 domains present in the carboxy end of the ArgBPs. One member of the Arg/Abl binding protein family, ArgBP2 is expressed in epithelial and cardiac muscle cells. The subcellular localization of ArgBP2 suggests that it functions as an adapter protein in the assembly of signaling complexes in stress fibers, and that it is a potential link between Abl family kinases and the actin cytoskeleton. The human ArgBP2 gene, which maps to chromosome 4, encodes multiple transcripts that yield different isoforms of ArgBP2. Another member of the ArgBP family, nArgBP2, which is specifically expressed in neural tissue, has the carboxy terminal SH3 domains characteristic of Arg/Abl binding proteins, as well as a sorbin homology domain near the N-terminus and a zinc finger motif in the middle region of the protein. nArgBP2 interacts with the proline rich region of SAPAP via its third SH3 domain. In rat brain, nArgBP2 colocalizes with SAPAP at the synapses of the cerebellum.

REFERENCES

1. Kruh, G.D., et al. 1990. The complete coding sequence of Arg defines the Abelson subfamily of cytoplasmic tyrosine kinases. *Proc. Natl. Acad. Sci. USA* 87: 5802-5806.
2. Wang, B., et al. 1997. ArgBP2, a multiple Src homology 3 domain-containing, Arg/Abl-interacting protein, is phospho-rylated in v-Abl-transformed cells and localized in stress fibers and cardiocyte Z-disks. *J. Biol. Chem.* 272: 17542-17550.
3. Kawabe, H., et al. 1999. nArgBP2, a novel neural member of ponsin/ArgBP2/vinexin family that interacts with synapse-associated protein 90/postsynaptic density-95-associated protein (SAPAP). *J. Biol. Chem.* 274: 30914-30918.
4. Hirao, K., et al. 2000. Association of synapse-associated protein 90/postsynaptic density-95-associated protein (SAPAP) with neurofilaments. *Genes Cells* 5: 203-210.

CHROMOSOMAL LOCATION

Genetic locus: Sorbs2 (mouse) mapping to 8 B1.1.

PRODUCT

ArgBP2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ArgBP2 shRNA Plasmid (m): sc-141193-SH and ArgBP2 shRNA (m) Lentiviral Particles: sc-141193-V as alternate gene silencing products.

For independent verification of ArgBP2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141193A, sc-141193B and sc-141193C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ArgBP2 siRNA (m) is recommended for the inhibition of ArgBP2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

ArgBP2 (B-11): sc-514671 is recommended as a control antibody for monitoring of ArgBP2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ArgBP2 gene expression knockdown using RT-PCR Primer: ArgBP2 (m)-PR: sc-141193-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.