

ARL5B siRNA (m): sc-141249

BACKGROUND

ADP-ribosylation factors (ARFs) are highly conserved guanine nucleotide binding proteins that enhance the ADP-ribosyltransferase activity of Cholera Toxin. ARFs are important in eukaryotic vesicular trafficking pathways, and they play an essential role in the activation of phospholipase D (PC-PLD). ARL5B (ADP-ribosylation factor-like protein 5B), also known as ARL8, is a 179 amino acid protein that belongs to the RAS superfamily of regulatory GTPases. Human ARL5B shares 100% identity with mouse Arl5b and 71% identity with the *Drosophila* homolog. Expressed in brain, heart, lung, cartilage and kidney, but not in spleen, ARL5B is most related to ARL5, with which it shares 80% protein sequence identity. Two isoforms of ARL5B exist as a result of alternative splicing events.

REFERENCES

1. Pasqualato, S., et al. 2002. ARF, ARL, ARP and SAR proteins: a family of GTP-binding proteins with a structural device for "front-back" communication. *EMBO Rep.* 3: 1035-1041.
2. Sebald, E., et al. 2003. Isolation of a new member of the ADP-ribosylation like factor gene family, ARL8, from a cartilage cDNA library. *Gene* 311: 147-151.
3. Okai, T., et al. 2004. Novel small GTPase subfamily capable of associating with tubulin is required for chromosome segregation. *J. Cell Sci.* 117: 4705-4715.
4. Burd, C.G., et al. 2004. ARF-like GTPases: not so ARF-like after all. *Trends Cell Biol.* 14: 687-694.
5. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 608909. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Kahn, R.A., et al. 2005. ARF family GTPases: roles in membrane traffic and microtubule dynamics. *Biochem. Soc. Trans.* 33: 1269-1272.
7. Haraguchi, T., et al. 2006. Expression of ADP-ribosylation factor-like protein 8B mRNA in the brain is down-regulated in mice fed a high-fat diet. *Biosci. Biotechnol. Biochem.* 70: 1798-1802.
8. Kahn, R.A., et al. 2006. Nomenclature for the human Arf family of GTP-binding proteins: ARF, ARL, and SAR proteins. *J. Cell Biol.* 172: 645-650.
9. Hofmann, I. and Munro, S. 2006. An N-terminally acetylated ARF-like GTPase is localised to lysosomes and affects their motility. *J. Cell Sci.* 119: 1494-1503.

CHROMOSOMAL LOCATION

Genetic locus: Arl5b (mouse) mapping to 2 A2.

PRODUCT

ARL5B siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ARL5B shRNA Plasmid (m): sc-141249-SH and ARL5B shRNA (m) Lentiviral Particles: sc-141249-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ARL5B siRNA (m) is recommended for the inhibition of ARL5B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

ARL5B (E-3): sc-393511 is recommended as a control antibody for monitoring of ARL5B gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ARL5B gene expression knockdown using RT-PCR Primer: ARL5B (m)-PR: sc-141249-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.