



# ATP5L siRNA (m): sc-141354

## BACKGROUND

Mitochondrial ATP synthase is composed of two multi-subunit complexes that utilize an inner membrane electrochemical gradient to catalyze the synthesis of ATP during oxidative phosphorylation. The two multi-subunit complexes are designated  $F_1$  and  $F_0$ , the former of which comprises the soluble catalytic core and the latter of which comprises the membrane-spanning proton channel of ATP synthase.  $F_1$  consists of five distinct subunits, designated ATP5A, ATP5B, ATP5C1, ATP5D and ATP5E, while  $F_0$  consists of ten subunits, designated ATP5H, ATP5G1, ATP5I, ATP5G2, ATP5J2, ATP5J, ATP5G3, ATP5S, ATP5F1 and ATP5L. ATP5L, also known as ATP5JG, is a 103 amino acid protein that localizes to the mitochondrial membrane and exists as a subunit of the  $F_0$  complex.

## REFERENCES

1. Elston, T., et al. 1998. Energy transduction in ATP synthase. *Nature* 391: 510-513.
2. Wang, H. and Oster, G. 1998. Energy transduction in the  $F_1$  motor of ATP synthase. *Nature* 396: 279-282.
3. Aggeler, R., et al. 2002. A functionally active human  $F_1F_0$  ATPase can be purified by immunocapture from heart tissue and fibroblast cell lines. Subunit structure and activity studies. *J. Biol. Chem.* 277: 33906-33912.
4. Leyva, J.A., et al. 2003. Understanding ATP synthesis: structure and mechanism of the  $F_1$ -ATPase (review). *Mol. Membr. Biol.* 20: 27-33.
5. Oster, G. and Wang, H. 2003. Rotary protein motors. *Trends Cell Biol.* 13: 114-121.
6. Cross, R.L. 2004. Molecular motors: turning the ATP motor. *Nature* 427: 407-408.

## CHROMOSOMAL LOCATION

Genetic locus: Atp5l (mouse) mapping to 9 A5.2.

## PRODUCT

ATP5L siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ATP5L shRNA Plasmid (m): sc-141354-SH and ATP5L shRNA (m) Lentiviral Particles: sc-141354-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^\circ\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^\circ\text{C}$ , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ATP5L siRNA (m) is recommended for the inhibition of ATP5L expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.